

Truncation errors: using Taylor series to approximation functions

Approximating functions using polynomials:

Let's say we want to approximate a function $f(x)$ with a polynomial

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + \dots$$

For simplicity, assume we know the function value and its derivatives at $x_0 = 0$ (we will later generalize this for any point). Hence,

$$f'(x) = a_1 + 2 a_2 x + 3 a_3 x^2 + 4 a_4 x^3 + \dots$$

$$f''(x) = 2 a_2 + (3 \times 2) a_3 x + (4 \times 3) a_4 x^2 + \dots$$

$$f'''(x) = (3 \times 2) a_3 + (4 \times 3 \times 2) a_4 x + \dots$$

$$f^{iv}(x) = (4 \times 3 \times 2) a_4 + \dots$$

$$f(0) = a_0 \quad f''(0) = 2 a_2 \quad f^{iv}(0) = (4 \times 3 \times 2) a_4$$

$$f'(0) = a_1 \quad f'''(0) = (3 \times 2) a_3$$

$$f^{(i)}(0) = i! a_i$$

Taylor Series

Taylor Series approximation about point $x_0 = 0$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + \dots$$

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

$$f(x) = \sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} x^i$$

Taylor Series

In a more general form, the Taylor Series approximation about point x_o is given by:

$$f(x) = f(x_o) + f'(x_o)(x - x_o) + \frac{f''(x_o)}{2!} (x - x_o)^2 + \frac{f'''(x_o)}{3!} (x - x_o)^3 + \dots$$

$$f(x) = \sum_{i=0}^{\infty} \frac{f^{(i)}(x_o)}{i!} (x - x_o)^i$$

Clicker question

Assume a finite Taylor series approximation that converges everywhere for a given function $f(x)$ and you are given the following information:

$$f(1) = 2; f'(1) = -3; f''(1) = 4; f^{(n)}(1) = 0 \forall n \geq 3$$

Evaluate $f(4)$

- A) 29
- B) 11
- C) -25
- D) -7
- E) None of the above

Taylor Series

We cannot sum infinite number of terms, and therefore we have to **truncate**.

How **big is the error** caused by truncation? Let's write $h = x - x_0$

$$f(x_0 + h) - \sum_{i=0}^n \frac{f^{(i)}(x_0)}{i!} (h)^i = \sum_{i=n+1}^{\infty} \frac{f^{(i)}(x_0)}{i!} (h)^i$$

And as $h \rightarrow 0$ we write:

$$\left| f(x_0 + h) - \sum_{i=0}^n \frac{f^{(i)}(x_0)}{i!} (h)^i \right| \leq C \cdot h^{n+1}$$

**Error due to Taylor
approximation of
degree n**

$$\left\{ \left| f(x_0 + h) - \sum_{i=0}^n \frac{f^{(i)}(x_0)}{i!} (h)^i \right| = O(h^{n+1}) \right.$$

Taylor series with remainder

Let f be $(n + 1)$ -times differentiable on the interval (x_o, x) with $f^{(n)}$ continuous on $[x_o, x]$, and $h = x - x_o$

$$f(x_o + h) - \sum_{i=0}^n \frac{f^{(i)}(x_o)}{i!} (h)^i = \sum_{i=n+1}^{\infty} \frac{f^{(i)}(x_o)}{i!} (h)^i$$

Then there exists a $\xi \in (x_o, x)$ so that

$$f(x_o + h) - \sum_{i=0}^n \frac{f^{(i)}(x_o)}{i!} (h)^i = \underbrace{\frac{f^{(n+1)}(\xi)}{(n+1)!} (\xi - x_o)^{n+1}}_{\text{Taylor remainder}} \quad f(x) - T(x) = R(x)$$

And since $|\xi - x_o| \leq h$

$$f(x_o + h) - \sum_{i=0}^n \frac{f^{(i)}(x_o)}{i!} (h)^i \leq \frac{f^{(n+1)}(\xi)}{(n+1)!} (h)^{n+1}$$



Demo: Polynomial Approximation with Derivatives

`f`

$$\sqrt{-x^2 + 1}$$

`taylor`

$$-\frac{x^2}{2} + 1$$



$$\text{error} = \text{taylor} - f$$

Demo: Polynomial Approximation with Derivatives



f

$$\sqrt{-x^2 + 1}$$

taylor

$$-\frac{x^2}{2} + 1$$

error = taylor - f



Clicker question

Error Order for Taylor series

1 point

The series expansion for e^x about 2 is

$$\exp(2) \cdot \left(1 + (x - 2) + \frac{(x - 2)^2}{2!} + \frac{(x - 2)^3}{3!} + \dots \right).$$

If we evaluate e^x using only the first four terms of this expansion (i.e. only terms up to and including $\frac{(x-2)^3}{3!}$), then what is the error in big-O notation?

Choice*

- A) $O(x^4)$
- B) $O(x^5)$
- C) $O(x^3)$
- D) $O((x - 2)^3)$
- E) $O((x - 2)^4)$

Demo “Taylor of $\exp(x)$ about 2”

Making error predictions

Suppose you expand $\sqrt{x - 10}$ in a Taylor polynomial of degree 3 about the center $x_0 = 12$. For $h_1 = 0.5$, you find that the Taylor truncation error is about 10^{-4} .

What is the Taylor truncation error for $h_2 = 0.25$?

$\text{Error}(h) = O(h^{n+1})$, where $n = 3$, i.e.

$$\text{Error}(h_1) \approx C \cdot h_1^4$$

$$\text{Error}(h_2) \approx C \cdot h_2^4$$

While not knowing C or lower order terms, we can use the ratio of h_2/h_1

$$\text{Error}(h_2) \approx C \cdot h_2^4 = C \cdot h_1^4 \left(\frac{h_2}{h_1}\right)^4 \approx \text{Error}(h_1) \cdot \left(\frac{h_2}{h_1}\right)^4$$

Can make prediction of the error for one h if we know another.

Using Taylor approximations to obtain derivatives

Let's say a function has the following Taylor series expansion about $x = 2$.


$$f(x) = \frac{5}{2} - \frac{5}{2}(x - 2)^2 + \frac{15}{8}(x - 2)^4 - \frac{5}{4}(x - 2)^6 + \frac{25}{32}(x - 2)^8 + O((x - 2)^9)$$

Therefore the Taylor polynomial of order 4 is given by

$$t(x) = \frac{5}{2} - \frac{5}{2}(x - 2)^2 + \frac{15}{8}(x - 2)^4$$

where the first derivative is

$$t'(x) = -5(x - 2) + \frac{15}{2}(x - 2)^3$$



Using Taylor approximations to obtain derivatives

We can get the approximation for the derivative of the function $f(x)$ using the derivative of the Taylor approximation:

$$t'(x) = -5(x - 2) + \frac{15}{2}(x - 2)^3$$


For example, the approximation for $f'(2.3)$ is

$$f'(2.3) \approx t'(2.3) = -1.2975$$

(note that the exact value is

$$f'(2.3) = -1.31444$$

What happens if we want to use the same method to approximate $f'(3)$?



Clicker question

The function

$$f(x) = \cos(x) x^2 + \frac{\sin(2x)}{(x+2x^2)^3}$$

is approximated by the following Taylor polynomial of degree $n = 2$ about $x = 2\pi$

$$t_2(x) = 39.4784 + 12.5664 (x - 2\pi) - 18.73922 (x - 2\pi)^2$$

Determine an approximation for the first derivative of $f(x)$ at $x = 6.1$

- A) 18.7741
- B) 12.6856
- C) 19.4319
- D) 15.6840

Computing integrals using Taylor Series

A function $f(x)$ is approximated by a Taylor polynomial of order n around $x = 0$.

$$t_n = \sum_{i=0}^n \frac{f^{(i)}(0)}{i!} (x)^i$$

We can find an approximation for the integral $\int_s^t f(x)dx$ by integrating the polynomial:

$$\begin{aligned} \int_s^t f(x)dx &\approx \int_s^t a_0 + a_1x + a_2x^2 + a_3x^3 dx \\ &= a_0 \int_s^t 1dx + a_1 \int_s^t x \cdot dx + a_2 \int_s^t x^2 dx + a_3 \int_s^t x^3 dx \end{aligned}$$

Where we can use $\int_s^t x^i dx = \frac{t^{i+1}}{i+1} - \frac{s^{i+1}}{i+1}$

Clicker question

A function $f(x)$ is approximated by the following Taylor polynomial:

$$t_5(x) = 10 + x - 5x^2 - \frac{x^3}{2} + \frac{5x^4}{12} + \frac{x^5}{24} - \frac{x^6}{72}$$

Determine an approximated value for $\int_{-3}^1 f(x) dx$

- A) -10.27
- B) -11.77
- C) 11.77
- D) 10.27