


1 Consider the following solitaire game. The puzzle consists of an $n \times m$ grid of squares, where each square may be empty, occupied by a red stone, or occupied by a blue stone. The goal of the puzzle is to remove some of the given stones so that the remaining stones satisfy two conditions:

- (1) every row contains at least one stone, and
- (2) no column contains stones of both colors.

For some initial configurations of stones, reaching this goal is impossible.



Prove that it is NP-hard to determine, given an initial configuration of red and blue stones, whether this puzzle can be solved.

Solution:

We show that this puzzle is NP-hard by describing a reduction from 3SAT.

Let Φ be a 3CNF boolean formula with m variables and n clauses. We transform this formula into a puzzle configuration in polynomial time as follows. The size of the board is $n \times m$. The stones are placed as follows, for all indices i and j :

- If the variable x_j appears in the i th clause of Φ , we place a blue stone at (i, j) .
- If the negated variable \bar{x}_j appears in the i th clause of Φ , we place a red stone at (i, j) .
- Otherwise, we leave cell (i, j) blank.

We claim that this puzzle has a solution if and only if Φ is satisfiable. This claim immediately implies that solving the puzzle is NP-hard. We prove our claim as follows:

- \implies First, suppose Φ is satisfiable; consider an arbitrary satisfying assignment. For each index j , remove stones from column j according to the value assigned to x_j :
- If $x_j = \text{TRUE}$, remove all red stones from column j .
 - If $x_j = \text{FALSE}$, remove all blue stones from column j .

In other words, remove precisely the stones that correspond to FALSE literals. Because every variable appears in at least one clause, each column now contains stones of only one color (if any). On the other hand, each clause of Φ must contain at least one TRUE literal, and thus each row still contains at least one stone. We conclude that the puzzle is satisfiable.

- ⇐ On the other hand, suppose the puzzle is solvable; consider an arbitrary solution. For each index j , assign a value to x_j depending on the colors of stones left in column j :
- If column j contains blue stones, set $x_j = \text{TRUE}$.
 - If column j contains red stones, set $x_j = \text{FALSE}$.
 - If column j is empty, set x_j arbitrarily.


In other words, assign values to the variables so that the literals corresponding to the remaining stones are all TRUE. Each row still has at least one stone, so each clause of Φ contains at least one TRUE literal, so this assignment makes $\Phi = \text{TRUE}$. We conclude that Φ is satisfiable.

This reduction clearly requires only polynomial time.

Rubric:[for all polynomial-time reductions] 10 points =

- + 3 points for the reduction itself
 - For an NP-hardness proof, the reduction must be from a known NP-hard problem. You can use any of the NP-hard problems listed in the lecture notes (except the one you are trying to prove NP-hard, of course).
- + 3 points for the “if” proof of correctness
- + 3 points for the “only if” proof of correctness
- + 1 point for writing “polynomial time”
- An incorrect polynomial-time reduction that still satisfies half of the correctness proof is worth at most 4/10.
- A reduction in the wrong direction is worth 0/10.


2 A *double-Hamiltonian tour* in an undirected graph G is a closed walk that visits every vertex in G exactly twice. Prove that it is NP-hard to decide whether a given graph G has a double-Hamiltonian tour.



This graph contains the double-Hamiltonian tour $a \rightarrow b \rightarrow d \rightarrow g \rightarrow e \rightarrow b \rightarrow d \rightarrow c \rightarrow f \rightarrow a \rightarrow c \rightarrow f \rightarrow g \rightarrow e \rightarrow a$.

Solution:

We prove the problem is NP-hard with a reduction from the standard Hamiltonian cycle problem. Let G be an arbitrary undirected graph. We construct a new graph H by attaching a small gadget to every vertex of G . Specifically, for each vertex v , we add two vertices $v^\#$ and v^b , along with three edges vv^b , $vv^\#$, and $v^bv^\#$.



A vertex in G , and the corresponding vertex gadget in H .

I claim that G has a Hamiltonian cycle if and only if H has a double-Hamiltonian tour.


\implies Suppose G has a Hamiltonian cycle $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_n \rightarrow v_1$. We can construct a double-Hamiltonian tour of H by replacing each vertex v_i with the following walk:

$$\dots \rightarrow v_i \rightarrow v_i^b \rightarrow v_i^\sharp \rightarrow v_i^b \rightarrow v_i^\sharp \rightarrow v_i \rightarrow \dots$$

\impliedby Conversely, suppose H has a double-Hamiltonian tour D . Consider any vertex v in the original graph G ; the tour D must visit v exactly twice. Those two visits split D into two closed walks, each of which visits v exactly once. Any walk from v^b or v^\sharp to any other vertex in H must pass through v . Thus, one of the two closed walks visits only the vertices v, v^b , and v^\sharp . Thus, if we simply remove the vertices in $H \setminus G$ from D , we obtain a closed walk in G that visits every vertex in G once.

Given any graph G , we can clearly construct the corresponding graph H in polynomial time.

With more effort, we can construct a graph H that contains a double-Hamiltonian tour *that traverses each edge of H at most once* if and only if G contains a Hamiltonian cycle. For each vertex v in G we attach a more complex gadget containing five vertices and eleven edges, as shown on the next page.




A vertex in G , and the corresponding modified vertex gadget in H .

Solution:

Bad and incorrect solution!!!

We attempt to prove the problem is NP-hard with a reduction from the Hamiltonian cycle problem. Let G be an arbitrary undirected graph. We construct a new graph H by attaching a self-loop every vertex of G . Given any graph G , we can clearly construct the corresponding graph H in polynomial time.




An incorrect vertex gadget.

Suppose G has a Hamiltonian cycle $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_n \rightarrow v_1$. We can construct a double-Hamiltonian tour of H by alternating between edges of the Hamiltonian cycle and self-loops:

$$v_1 \rightarrow v_1 \rightarrow v_2 \rightarrow v_2 \rightarrow v_3 \rightarrow \dots \rightarrow v_n \rightarrow v_n \rightarrow v_1.$$

On the other hand, if H has a double-Hamiltonian tour, we *cannot* conclude that G has a Hamiltonian cycle, because we cannot guarantee that a double-Hamiltonian tour in H uses *any* self-loops. The graph G shown below is a counterexample; it has a double-Hamiltonian tour (even before adding self-loops) but no Hamiltonian cycle.



This graph has a double-Hamiltonian tour.

Rubric:[for all polynomial-time reductions] 10 points =

- + 3 points for the reduction itself
 - For an NP-hardness proof, the reduction must be from a known NP-hard problem. You can use any of the NP-hard problems listed in the lecture notes (except the one you are trying to prove NP-hard, of course).
- + 3 points for the “if” proof of correctness
- + 3 points for the “only if” proof of correctness
- + 1 point for writing “polynomial time”
- An incorrect polynomial-time reduction that still satisfies half of the correctness proof is worth at most 4/10.
- A reduction in the wrong direction is worth 0/10.