


Lecture 20

Phase Transitions


- Phase diagrams
- Latent heats
- Phase-transition fun

Reading for this Lecture:
Elements Ch 13


Solids, Liquids and Gases



Solid



Liquid



Gas

Solids have fixed relationships among the atoms (or molecules)

Liquids have looser relationships among atoms.

⇒ The liquid has more entropy, because there are more ways to arrange atoms in the liquid.

In liquids there are still some correlations between atoms, but in gases there are essentially none. In most situations the entropy of the gas is vastly larger than that of the liquid or solid.

Phase Transitions and Entropy

If $S_{\text{gas}} > S_{\text{liquid}} > S_{\text{solid}}$, why are different phases stable at different temperatures?

The answer is that we must also consider the entropy of the environment. That's what free energy and chemical potential do for us.



For example: At low enough temperatures a substance like water is a solid. Its entropy is lower than that of the liquid so it must give up enough energy to its environment to make the total entropy increase when ice forms:

$$\text{Liquid H}_2\text{O} \rightarrow \text{Solid: } \Delta S_{\text{tot}} = \underbrace{\Delta S_{\text{L}} + \Delta S_{\text{S}}}_{< 0} + \underbrace{\Delta S_{\text{env}}}_{> 0} \geq 0$$

In order for this to work, enough heat must be given to the environment to make $\Delta S_{\text{tot}} \geq 0$.

OK. So, why is liquid H_2O favored at higher temperatures?

The relative sizes of the ΔS terms must be different.



Let's look at the problem more quantitatively.

Solid-gas equilibrium: vapor pressure


Consider solid-gas equilibrium at constant volume and temperature. Some substances (e.g., CO₂) don't exist as liquids at atmospheric pressure.

The solid has negligible entropy (compared to the gas), so we'll ignore it. In that case:

$$F_s = U_s - TS_s = -N\Delta \Rightarrow \mu_s = -\Delta$$

Binding energy of an atom in the solid

$$\text{The gas: } \mu_g = kT \ln(n/n_Q)$$



Set them equal and solve for the equilibrium gas pressure:

$$kT \ln\left(\frac{n}{n_Q}\right) = kT \ln\left(\frac{p}{p_Q}\right) = -\Delta \Rightarrow p \equiv p_{\text{vapor}} = p_Q e^{-\Delta/kT}$$

The equilibrium pressure is called the “vapor pressure” of the solid at temperature T. If $p < p_{\text{vapor}}$, atoms will leave the solid until $p_{\text{gas}} = p_{\text{vapor}}$. This is called sublimation. For liquids, it's called evaporation.

Examples: Si (28 g/mol): $p_{\text{vapor}} = (4.04 \times 10^4 \text{ atm})(28)^{3/2} e^{-3\text{eV}/.026\text{eV}} = 5 \times 10^{-44} \text{ atm}$
 CO₂ (44 g/mol): $p_{\text{vapor}} = (4.04 \times 10^4 \text{ atm})(44)^{3/2} e^{-0.26\text{eV}/.026\text{eV}} = 535 \text{ atm}$

Some solids don't sublime. Some do.
 Question: Does water ice sublime?

↑
 Note the different Δ's

Chemical Potential of an Ideal Gas

Remember: $\mu_g = kT \ln \frac{n}{n_Q} = kT \ln \frac{p}{p_Q}$, where $p_Q = n_Q kT$


It's negative, because (unless highly compressed) $n_Q \gg n$ and $p_Q \gg p$.

For helium at $T = 300\text{K}$ and $p = 1\text{ atm}$: $p_Q = (4.04 \times 10^4\text{ atm})(4)^{1.5} = 3.23 \times 10^5\text{ atm}$

So, $\mu_{\text{He}} = kT \ln \frac{p}{p_Q} = kT \ln(3.10 \times 10^{-6}) = (.026\text{ eV})(-12.69) = -0.33\text{ eV}$

All ideal gases will behave similarly, with a logarithmic pressure dependence and an approximately linear temperature dependence. (The curvature is due to the T dependence of n_Q .)

Here's a graph:




Vapor Pressure of a Solid and the p-T Phase Diagram

We saw that μ_{solid} is essentially constant ($-\Delta$). We can use this to determine graphically the equilibrium pressure (vapor pressure) of the solid as a function of T . For a given temperature, at what pressure does the curve cross $\mu = -\Delta$?

From this, we can plot the equilibrium curve, $p(T)$, where the two phases can coexist. This graph is called a p-T phase diagram. It tells us what regions in the (p, T) plane make a gas and which make a solid. The equilibrium curve separates them.

It's just a way to visualize the equilibrium equation:

$$\mu_g = \mu_s \rightarrow kT \ln \frac{p}{p_Q} = -\Delta \rightarrow p = p_Q e^{-\Delta/kT}$$



Solids and Liquids, More Accurately

We ignored the entropy of the solid, because the entropy of the gas was so much larger. **When comparing solids and liquids, we can't ignore entropy.** The entropy is small, but not completely negligible.

Compare solids and liquids (for a given substance):

- $\Delta_S > \Delta_L$. Atoms in the solid are more strongly bound.
- $S_S < S_L$. Atoms in the liquid have more available microstates (due to motion).


Compare the chemical potentials:

Remember that $F = U - TS$, and μ is the free energy per particle.

- At very low T , the entropy is not important, so $\mu_S < \mu_L$.
- As T increases, both chemical potentials decrease, but the liquid decreases faster.

Therefore, the solid phase is stable at low temperature, while the liquid phase is stable at high temperatures.


There is an equilibrium temperature, the melting/freezing temperature. As the substance passes through T_{freeze} , there is a phase transition.



Act 1: Solid-Liquid

1) Which point corresponds to a liquid?

A B C



2) The substance is in **state C**. What will happen?

- A) The substance will melt
- B) Free energy will minimize itself
- C) Entropy will maximize

Solution

1) Which point corresponds to a liquid?

A **B** C

Both points lie on the liquid's $\mu(T)$ curve.

Point A does not lie on the solid's curve, so it must be liquid

(but not one in equilibrium [lowest possible μ]).


At point B the liquid and solid are in equilibrium.

We don't actually know how much of each there is.

FYI: Global equilibrium condition:

$$\text{Minimum } G = N_g\mu_g + N_s\mu_s + N_l\mu_l \quad G = \text{Gibbs Free Energy}$$

Except at a coexistence temperature, T_c , two of the N 's will be zero in equilibrium.



2) The substance is in **state C**. What will happen?

- A) The substance will melt
- B) Free energy will minimize itself
- C) Entropy will maximize

Solution

1) Which point corresponds to a liquid?

A **B** C


Both points lie on the liquid's $\mu(T)$ curve.

Point A does not lie on the solid's curve, so it must be liquid

(but not one in equilibrium [lowest possible μ]).

At point B the liquid and solid are in equilibrium.

We don't actually know how much of each there is.



FYI: Global equilibrium condition:

Minimum $G = N_g\mu_g + N_s\mu_s + N_l\mu_l$ $G =$ Gibbs Free Energy

Except at a coexistence temperature, T_c , two of the N 's will be zero in equilibrium.



2) The substance is in **state C**. What will happen?

- A) The substance will melt**
- B) Free energy will minimize itself**
- C) Entropy will maximize**

B and C are equivalent. That's what systems do unless we interfere.

A is also correct, because $\mu_L < \mu_S$ at that temperature.

The p-T Diagram with Three Phases




At a given p, T , the system will reside in the lowest μ .

Phase transitions occur where the $\mu(T)$ curves cross.

If we raise T at constant p , the sequence of phase transitions depends on the value of p .

The solid and liquid curves don't change with pressure, because solids and liquids are nearly incompressible.

The p-T Diagram with Three Phases




At low pressure, the liquid phase is not stable for any T . The substance sublimates at temperature T_1 .

If we increase the pressure, the gas curve moves to the right. There is a critical pressure for which all three curves pass through a point. The three phases can coexist only at (T_2, p_2) .

At high pressure, there are two phase transitions: solid-liquid at T_3 , and liquid-gas at T_4 .

Phase Diagram:




Phase Transitions: OOPS!

The photos below show the results of steam cleaning a railroad tank car, then sealing it before the steam had cooled



Phase Diagrams



- On special transition lines, two phases are stable.
- At very special “triple points”, those line cross and three phases are stable.
- Notice that in almost all the (p.T) plane, only ONE phase is stable.


ACT 2: Phase Transitions

Let's think a bit about energy flow during phase transitions.

Suppose heat flows slowly into 0°C water containing some ice cubes (maintaining equilibrium).

What happens to the energy added to the system?

- A) It warms the ice.
- B) It warms the water.
- C) It breaks ice bonds.



Solution


Let's think a bit about energy flow during phase transitions.

Suppose heat flows slowly into 0°C water containing some ice cubes (maintaining equilibrium).

What happens to the energy added to the system?



- A) It warms the ice.
- B) It warms the water.
- C) It breaks ice bonds.

The temperature will remain constant until all the ice is melted.



Latent Heat

Consider a liquid-gas phase transition at constant pressure. The phase diagram doesn't tell the whole story. Look at how the system changes as we add heat:




Plot temperature vs the amount of heat added:

Remember the 1st law: $Q = \Delta U + p\Delta V$. Some of the energy goes into breaking the bonds (ΔU), and some goes into raising the weight ($p\Delta V$).

The p-T diagram doesn't show either of these changes.

The heat we need to add to effect the phase change is called "latent heat".



Heat of transformation
≡ Latent Heat

Latent Heat (2)

The heat required to convert the liquid entirely into gas is directly related to the entropy change of the material:

$$Q_{12} = T_1 \Delta S_{12} = \Delta U + p\Delta V \equiv \Delta H_{lg}$$

$H = U + pV$ is defined as the **Enthalpy** ('heat content') of a material.


For phase transitions:

“Heat of vaporization” = “enthalpy change” = “latent heat”:

$$\Delta H_{lg} = H_{\text{gas}} - H_{\text{liquid}} = \text{Latent Heat} = Q_L$$

Latent heats are typically given in units of J/mol or J/gram

Latent Heats of Evaporation			
Gas	Boiling temp (K)	ΔH (J/mol)	ΔS (J/mol·K)
Helium	4.2	92	22
Nitrogen	77	5,600	72
Oxygen	90	6,800	76
H ₂ O	370	40,000	120



There is a similar latent heat for the solid-liquid transition.

For H₂O at 1 atm:

$$Q_L(\text{ice-liquid}) = 80 \text{ cal/g} = 333 \text{ kJ/kg}$$

$$Q_L(\text{liquid-steam}) = 540 \text{ cal/g} = 2256 \text{ kJ/kg}$$