

Useful formulae

Momentum of a particle:

$$\vec{p} = m \frac{d\vec{r}}{dt}$$

Force on a particle:

$$\vec{F} = \frac{d\vec{p}}{dt}$$

Center of mass:

$$\vec{r}_{cm} = \frac{1}{M} \int \vec{r} dm = \frac{1}{M} \sum_i m_i \vec{r}_i \quad (1)$$

Change in momentum of center of mass in presence of external forces F_i :

$$\frac{d\vec{p}_{cm}}{dt} = \sum_i \vec{F}_i \quad (2)$$

Cylindrical coordinates:

$$x = R \cos \phi \quad (3)$$

$$y = R \sin \phi \quad (4)$$

$$z = z \quad (5)$$

$$R = \sqrt{x^2 + y^2} \quad (6)$$

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (7)$$

$$\hat{\mathbf{R}} = \cos \phi \hat{i} + \sin \phi \hat{j} \quad (8)$$

$$\hat{\phi} = -\sin \phi \hat{i} + \cos \phi \hat{j} \quad (9)$$

$$\frac{d\hat{\mathbf{R}}}{dt} = \dot{\phi} \hat{\phi} \quad (10)$$

$$\frac{d\hat{\phi}}{dt} = -\dot{\phi} \hat{\mathbf{R}} \quad (11)$$

$$\vec{v} = \dot{R} \hat{\mathbf{R}} + R \dot{\phi} \hat{\phi} + \dot{z} \hat{\mathbf{z}} \quad (12)$$

$$\vec{a} = (\ddot{R} - R \dot{\phi}^2) \hat{\mathbf{R}} + (R \ddot{\phi} + 2\dot{R}\dot{\phi}) \hat{\phi} + \ddot{z} \hat{\mathbf{z}} \quad (13)$$

$$\nabla^2 f = \frac{1}{R} \frac{\partial}{\partial R} (R \frac{\partial f}{\partial R}) + \frac{1}{R^2} \frac{\partial^2 f}{\partial \phi^2} + \frac{\partial^2 f}{\partial z^2} \quad (14)$$

Spherical coordinates (3D):

$$x = r \sin \theta \cos \phi \quad (15)$$

$$y = r \sin \theta \sin \phi \quad (16)$$

$$z = r \cos \theta \quad (17)$$

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (18)$$

$$\hat{\theta} = \cos \theta \cos \phi \hat{i} + \cos \theta \sin \phi \hat{j} - \sin \theta \hat{k} \quad (19)$$

$$\hat{\phi} = -\sin \phi \hat{i} + \cos \phi \hat{j} \quad (20)$$

$$\frac{d\hat{\mathbf{r}}}{dt} = \dot{\theta} \hat{\theta} + \sin \theta \dot{\phi} \hat{\phi} \quad (21)$$

$$\frac{d\hat{\theta}}{dt} = -\dot{\theta} \hat{\mathbf{r}} + \cos \theta \dot{\phi} \hat{\phi} \quad (22)$$

$$\frac{d\hat{\phi}}{dt} = -\dot{\phi} (\sin \theta \hat{\mathbf{r}} + \cos \theta \hat{\theta}) \quad (23)$$

$$\vec{v} = \dot{r} \hat{\mathbf{r}} + r \dot{\theta} \hat{\theta} + r \dot{\phi} \sin \theta \hat{\phi} \quad (24)$$

$$\begin{aligned} \vec{a} = & (\ddot{r} - r \dot{\theta}^2 - r \dot{\phi}^2 \sin^2 \theta) \hat{\mathbf{r}} + \\ & (r \ddot{\theta} + 2\dot{r}\dot{\theta} - r \dot{\phi}^2 \sin \theta \cos \theta) \hat{\theta} + \\ & (r \ddot{\phi} \sin \theta + 2\dot{r}\dot{\phi} \sin \theta + 2r\dot{\theta}\dot{\phi} \cos \theta) \hat{\phi} \end{aligned} \quad (25)$$

$$\begin{aligned} \nabla^2 f = & \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial f}{\partial r}) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta \frac{\partial f}{\partial \theta}) + \\ & \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2} \end{aligned} \quad (26)$$

Force from a potential:

$$\vec{F} = -\vec{\nabla} U \quad (27)$$

$$U(x) = - \int_{x_0}^x dx' F(x') \quad (28)$$

$$\frac{1}{2} m(v_2^2 - v_1^2) = \int_{x_1}^{x_2} dx' F(x') \quad (29)$$

Gradient of a scalar field:

$$\vec{\nabla} U = \frac{\partial U}{\partial x} \hat{i} + \frac{\partial U}{\partial y} \hat{j} + \frac{\partial U}{\partial z} \hat{k} \quad (30)$$

Curl of a vector field:

$$\vec{\nabla} \times \vec{F} = \hat{i} \left(\frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z} \right) + \hat{j} \left(\frac{\partial F_x}{\partial z} - \frac{\partial F_z}{\partial x} \right) + \hat{k} \left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) \quad (31)$$

Stokes' Theorem, which turns a line integral around a closed path into an integral over the enclosed area, with $d\vec{A}$ a vector pointing in the direction orthogonal to the area (for example, a

path that lies in the $x-y$ plane gets turned into an area integral with $d\vec{A}$ pointing in the z direction, which picks out the z component of the curl:

$$\oint \vec{F} \cdot d\vec{r} = \int (\vec{\nabla} \times \vec{F}) \cdot d\vec{A} \quad (32)$$

Simple harmonic motion:

$$U(x) = \frac{1}{2}k(x - x_0)^2 + \text{constant} \quad (33)$$

$$F(x) = -k(x - x_0) \quad (34)$$

$$\omega \equiv \sqrt{\frac{k}{m}} \quad (35)$$

$$\text{If } \ddot{x} = -\omega^2 x \text{ then } x = A \sin(\omega t + \phi) \quad (36)$$

Taylor expansion:

$$\begin{aligned} f(x) \sim & f(x_0) + \frac{df}{dx}(x_0)(x - x_0) \\ & + \frac{1}{2!} \frac{d^2f}{dx^2}(x_0)(x - x_0)^2 \\ & + \frac{1}{3!} \frac{d^3f}{dx^3}(x_0)(x - x_0)^3 + \dots \end{aligned} \quad (37)$$

Rocket-related useful equation (for exhaust relative velocity \vec{u} ; careful with signs!):

$$\frac{d\vec{P}}{dt} = M \frac{d\vec{v}}{dt} + \vec{u} \frac{dm}{dt} \quad (38)$$

Center of mass frame scattering (relative velocity \vec{V}):

$$\vec{V} = \vec{v}_1 - \vec{v}_2 \quad (39)$$

$$\vec{p}_1 = \mu \vec{V} = -\vec{p}_2 \quad (40)$$

$$\mu = \frac{m_1 m_2}{m_1 + m_2} \quad (41)$$

Potential energy:

$$\text{Constant force : } U(x) = -Fx + C \quad (42)$$

$$\text{Spring : } U(x) = \frac{1}{2}kx^2 + C \quad (43)$$

$$\text{1/r}^2 \text{ Gravity : } U(x) = \frac{-GMm}{r} + C \quad (44)$$

Some useful approximations (for small arguments to first order):

$$\sin \theta \sim \theta - \dots \quad (45)$$

$$\cos \theta \sim 1 - + \dots \quad (46)$$

$$\ln(1 + x) \sim x2 + \dots \quad (47)$$

$$e^x \sim 1 + x + \dots \quad (48)$$

$$(1 + x)^\alpha \sim 1 + \alpha x + \dots \quad (49)$$

Central force motion effective potential for reduced mass μ and angular momentum ℓ and potential $U(r)$:

$$U_{eff}(r) = U(r) + \frac{\ell^2}{2\mu r^2} \quad (50)$$

Orbit in a potential $U = -C/r$:

$$r(\theta) = \frac{(\ell^2/\mu C)}{1 - \sqrt{1 + (2E\ell^2/\mu C^2)}} \sin(\theta - \theta_0) \quad (51)$$

Taking $\theta_0 = -\pi/2$, and using the following definitions:

$$r = \frac{r_0}{1 - \epsilon \cos \theta} \quad (52)$$

$$\epsilon \equiv \sqrt{1 + \frac{2E\ell^2}{\mu C^2}} \quad (53)$$

$$r_0 \equiv \frac{\ell^2}{\mu C} \quad (54)$$

For a bound orbit, this is an ellipse, with the length of the longest axis A (all the way across i.e., the thing that would be the diameter, not the radius, for a circle) and the short axis B:

$$A = \frac{2r_0}{1 - \epsilon^2} = \frac{C}{-E} \quad (55)$$

$$B = \frac{2\ell}{\sqrt{-2\mu E}} \quad (56)$$

The longest and shortest distances from the central object:

$$r_{max} = \frac{r_0}{1 - \epsilon} \quad (57)$$

$$r_{min} = \frac{r_0}{1 + \epsilon} \quad (58)$$

Kepler's third law relating period T to length of long axis A :

$$T^2 = \frac{\pi^2}{2(M+m)G} A^3 \quad (59)$$

Vis-viva equation for Keplerian orbits:

$$v^2 = GM_{tot} \left(\frac{2}{r} - \frac{1}{a} \right) \quad (60)$$

Poisson's equation:

$$\nabla^2 \Phi = 4\pi G \rho \quad (61)$$

Fourier series:

$$f(t) = \frac{a_\circ}{2} + \sum_{n=1}^{\infty} [a_n \cos(n\omega t) + b_n \sin(n\omega t)] \quad (62)$$

$$\text{where } \omega \equiv \frac{2\pi}{T} \quad (63)$$

$$a_n = \frac{2}{T} \int_0^T f(t) \cos(n\omega t) dt \quad (64)$$

$$b_n = \frac{2}{T} \int_0^T f(t) \sin(n\omega t) dt \quad (65)$$

Fourier transforms:

$$\tilde{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt \quad (66)$$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \tilde{f}(\omega) e^{+i\omega t} d\omega \quad (67)$$

Lightly damped harmonic motion (for ϕ and A set by initial conditions):

$$\ddot{x} = -\gamma \dot{x} - \omega_\circ^2 x \quad (68)$$

$$x(t) = Ae^{-\gamma t/2} \cos(\omega' t + \phi) \quad (69)$$

$$\omega' \equiv \sqrt{\omega_\circ^2 - \frac{\gamma^2}{4}} \quad (70)$$

Forced damped harmonic motion: steady state solution (initial conditions are irrelevant

after a sufficiently long time):

$$\ddot{x} = -\gamma \dot{x} - \omega_\circ^2 x + \frac{F_\circ}{m} \cos(\omega_F t) \quad (71)$$

$$x_{ss}(t) = \frac{F_\circ \cos(\omega_F t - \phi)}{m \sqrt{(\omega_\circ^2 - \omega_F^2)^2 + \omega_F^2 \gamma^2}} \quad (72)$$

$$\tan(\phi) = \frac{\omega_F \gamma}{\omega_\circ^2 - \omega_F^2} \quad (73)$$

Green's function for damped oscillator, where $H(x) = 0$ for $x < 0$ and $H(x) = 1$ for $x > 0$:

$$G(t - \tau) = \frac{\sin[\omega'(t - \tau)]}{m\omega'} e^{-\gamma(t-\tau)/2} H(t - \tau) \quad (74)$$

$$|\tilde{G}(\omega)| = \frac{1}{m} \frac{1}{\sqrt{(\omega_\circ^2 - \omega^2)^2 + \gamma^2 \omega^2}} \quad (75)$$

$$x(t) = \int_{-\infty}^t F(\tau) G(t - \tau) d\tau \quad (76)$$

Fictitious force: uniform acceleration \vec{a} for mass m :

$$\vec{F}_{fict} = -m\vec{a} \quad (77)$$

Coriolis force, for system rotating with angular velocity Ω and for a velocity measured in the rotating frame of \vec{v}_{rot} :

$$\vec{F}_{cor} = -2m\vec{\Omega} \times \vec{v}_{rot} \quad (78)$$

Centrifugal force, for system rotating with angular velocity Ω :

$$\vec{F}_{cent} = -m\vec{\Omega} \times (\vec{\Omega} \times \vec{r}) \quad (79)$$

Relation between velocities measured in rotating and inertial frames:

$$\left(\frac{d\vec{B}}{dt} \right)_{inertial} = \left(\frac{d\vec{B}}{dt} \right)_{rotating} + \vec{\Omega} \times \vec{B} \quad (80)$$

Center of mass:

$$\vec{r}_{cm} = \frac{1}{M} \int \vec{r} dm = \frac{1}{M} \sum_i m_i \vec{r}_i \quad (81)$$

Change in momentum of center of mass in presence of external forces F_i :

$$\frac{d\vec{p}_{cm}}{dt} = \sum_i \vec{F}_i \quad (82)$$

Moment of inertia:

$$I_{zz} = \int (x^2 + y^2) dm \quad (83)$$

Angular momentum of a particle (with angular velocity $\omega = d\phi/dt$):

$$\vec{L} = \vec{r} \times \vec{p} = I\vec{\omega}$$

Torque on a particle:

$$\vec{\tau} = \vec{r} \times \vec{F} = I \frac{d\vec{\omega}}{dt}$$

Some useful moments of inertia (through center of mass for the direction that is the symmetry axis):

$$\text{Solid disk : } I = \frac{1}{2}MR^2 \quad (84)$$

$$\text{Ring : } I = MR^2 \quad (85)$$

$$\text{Solid sphere : } I = \frac{2}{5}MR^2 \quad (86)$$

Parallel Axis Theorem (moment of inertia for axis parallel to one through center of mass):

$$I = I_{cm} + Ml^2 \quad (87)$$

Kinetic energy of object with velocity of center of mass v_{cm} and rotating around c.o.m with angular velocity ω :

$$K = \frac{1}{2}mv_{cm}^2 + \frac{1}{2}I\omega^2 \quad (88)$$

Lagrangian Mechanics:

$$S = \int_{t_1}^{t_2} L dt \quad (89)$$

$$L = T - V \quad (90)$$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_\alpha} \right) = \frac{\partial L}{\partial q_\alpha} \quad (91)$$

In the presence of constraints $C(q_\alpha, t) = 0$, those may be handled as Lagrange multipliers, adding $\lambda C(q_\alpha, t)$ to the Lagrangian and proceeding as before.

A few trig identities:

$$\cos(a + b) = \cos a \cos b - \sin a \sin b \quad (92)$$

$$\sin(a + b) = \sin a \cos b + \cos a \sin b \quad (93)$$

Some more calculus:

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \dot{\vec{v}} \cdot \nabla f \quad (94)$$

$$\ddot{x} = \frac{1}{2} \frac{d}{dx} \dot{x}^2 \quad (95)$$

Some integrals:

$$\int \frac{dx}{x+a} = \ln(x+a) + C \quad (96)$$

$$\int e^x dx = e^x + C \quad (97)$$

$$\int \ln x dx = x \ln x - x + C \quad (98)$$

$$\int \sin x dx = -\cos x + C \quad (99)$$

$$\int \cos x dx = \sin x + C \quad (100)$$

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) \quad (101)$$

$$\int \frac{xdx}{\sqrt{x^2 + a^2}} = \sqrt{a^2 + x^2} \quad (102)$$

$$\int \frac{dx}{(a+x)^n} = \frac{(a+x)^{1-n}}{1-n} + C \quad (103)$$

$$\int \frac{xdx}{(x+a)^3} = -\frac{a+2x}{2(a+x)^2} \quad (104)$$

$$\int_{-\infty}^{\infty} f(t) \delta(t-a) dt = f(a) \quad (105)$$

$$\int_{-\infty}^{\infty} f(t) \delta(a-t) dt = f(a) \quad (106)$$

Some Fourier transform pairs:

$$\int_{-\infty}^{\infty} e^{-i\omega t} e^{iat} dt = 2\pi\delta(\omega - a) \quad (107)$$

$$\int_{-\infty}^{\infty} e^{-i\omega t} e^{-at^2} dt = \sqrt{\frac{\pi}{a}} e^{-\omega^2/(4a)} \quad (108)$$

$$\int_{-\infty}^{\infty} e^{-i\omega t} e^{-a|t|} dt = \frac{2a}{a^2 + \omega^2} \quad (109)$$

$$\int_{-\infty}^{\infty} e^{-i\omega t} \cos(at^2) dt = \sqrt{\frac{\pi}{a}} \cos\left(\frac{\omega^2}{4a} - \frac{\pi}{4}\right) \quad (110)$$

$$\int_{-\infty}^{\infty} e^{-i\omega t} \sin(at^2) dt = -\sqrt{\frac{\pi}{a}} \sin\left(\frac{\omega^2}{4a} - \frac{\pi}{4}\right) \quad (111)$$