
Millikan Oil Drop Experiment

Physics 401, Fall 2015
Eugene V. Colla





Agenda

- 1. Measuring of the charge of electron.
- 2. Robert Millikan and his oil drop experiment
- 3. Theory of the experiment
- 4. Laboratory setup
- 5. Data analysis



Measuring of the charge of the electron

- 1. Oil drop experiment. Robert A. Millikan.. (1909). $e=1.5924(17)\times 10^{-19}$ C
- 2. Shot noise experiment. First proposed by Walter H. Schottky
- 3. In terms of the Avogadro constant and Faraday constant e=

 $\frac{F}{N_A}$; F- Faraday constant, N_A - Avagadro constant. Best uncertainty ~1.6 ppm.

- 4. From Josephson $(K_J = \frac{2e}{h})$ and von Klitzing $\left(R_K = \frac{h}{e^2}\right)$ constants
- 5. Recommended by NIST value 1.602 176 565(35) 10⁻¹⁹ C

Robert Millikan. Oil Drop Experiment



ROBERT ANDREWS MILLIKAN 1868-1953



The Nobel Prize in Physics 1923. Robert A. Millikan "for his work on the elementary charge of electricity and on the photoelectric effect".

ON THE ELEMENTARY ELECTRICAL CHARGE AND THE

ELECTRICAL CHARGE AND AVOGADRO CONSTANT.

AVOGADRO CONSTANT.

By R. A. MILLIKAN.


I. Introductory.



University of Chicago 10/12/2016



Robert Millikan. Oil drop experiment



ROBERT ANDREWS MILLIKAN 1868-1953





Diagram and picture of apparatus





Oil Drop Experiment.

Motivation:

Measurement of the magnitude of the electron charge!

Demonstrate that the electron charge is quantized!






Measure the charge of an electron to ±3%

Picture of the PASCO setup



Oil Drop Experiment.




Forces on the oil drop:

- 1) Gravity + buoyant force (air displaced by oil drop)
- 2) Drag force of the oil drop in the air



Oil Drop Experiment.




Forces on the oil drop:


- 1) Gravity + buoyant force (air displaced by oil drop)
- 2) Drag force of the oil drop in the air
- 3) Electric force on oil drops which carry charge Q



Apparatus. Schematic Layout



Apparatus: Actual Setup



What is Measured

rise time measurement stops here

rise time t_{rise}

rise time measurement starts here

Allow drop to "undershoot" here before starting next rise time experiment

fall time measurement starts here

fall time $oldsymbol{t}_{oldsymbol{g}}$

x =fall distance = rise distance. x must be the same for all drops!

fall time measurement stops here



Balance of Forces: Newton's Law

$$\vec{E} \qquad \qquad \qquad \qquad \vec{F}_E = Q\vec{E}$$

$$\vec{F}_{drag} = -6\pi\eta \vec{a}\vec{v} \qquad \qquad \vec{F}_g = -mg\hat{z}$$

a: radius of drop

 ρ : density $\rho = \rho_{oil} - \rho_{air}$

v: velocity of oil drop

Q: charge of oil drop

E: electric field E=V/d

V : Voltage across plates

n: viscosity of air

: gravitational const.

Particle reached terminal velocity

$$\frac{d\vec{v}}{dt} = 0$$

$$|\vec{F}_g + \vec{F}_{drag} + \vec{F}_E| = 0$$

$$\vec{F}_{g} = -mg\hat{z} \tag{1}$$

$$\vec{F}_{g} = -mg\hat{z} \qquad (1)$$

$$\vec{F}_{drag} = -6\pi\eta a\vec{v} \quad (2)$$

$$\vec{F}_E = Q\vec{E} \tag{3}$$


$$\vec{F} = m \frac{d\vec{v}}{dt} = \vec{F}_g + \vec{F}_{drag} + \vec{F}_E$$

Forces on the oil drop:

- (1) Gravity + buoyant force (air displaced by oil drop)
- (2) Drag force of the oil drop in the air
- (3) Electric force on oil drops which carry charge Q

1μ size particle reaches the terminal velocity in ~10⁻⁵s

Modification to Stokes Law



George Gabriel Stokes (1819-1903)

$$\vec{F}_{drag} = -6\pi\eta a\vec{v}$$

For small particle radius (a<15 μ) Stokes law need to be corrected. This correction was derived by E. Cunningham.

$$\vec{F}_{drag} = -6\pi\eta \frac{a}{f_c} \vec{v}$$



Ebenezer Cunningham (1881-1977)

$$f_c = 1 + A \frac{\lambda}{a} + B \frac{\lambda}{a} e^{-c\frac{a}{\lambda}}, \quad A = 1.246, B = 0.42, C = 0.78$$

$$f_c \approx 1 + A \frac{\lambda}{a} = 1 + \frac{r_c}{a} \approx 1.1, \text{ for } a \approx 10^{-6} \text{m}, \quad r_c = \frac{6.18 \times 10^{-5}}{p[\text{mmHg}]}$$

Here a – particle radius; λ – mean free path of the gas molecules

negligible term

$$\lambda \text{ [m]} = 6.53 \times 10^{-8} \frac{760}{p \text{[mmHg]}}$$





rise time measurement stops here

rise time t_{rise}

rise time measurement starts here

Allow drop to "undershoot" here before starting next rise time experiment

fall time measurement starts here

fall time $oldsymbol{t_g}$

x =fall distance = rise distance. x must be the same for all drops!

fall time measurement stops here



Solving Newton's Law: Q(t_g, t_{rise})

 f_c can be found from Newton law equation in the case of V=0 (falling drop)

$$\vec{F}_g + \vec{F}_{drag} = \frac{4}{3} a^3 \rho g - 6\pi \eta \frac{a}{f_c} \vec{v} = 0$$

(see write-up)

$$\frac{1}{f_c^{\frac{2}{3}}} \approx 1 - \left(\frac{t_g}{\tau_g}\right)^{\frac{1}{2}} ; \quad \tau_g = \frac{2\eta x}{\rho g r_c^2}; \quad r_c[m] = \frac{6.18 \times 10^{-5}}{\rho [mmHg]}$$

Solving Newton's law: Q(t_g, t_{rise})

$$Q = n \times e = \frac{1}{f_c^{3/2}} \frac{9\pi d}{V} \sqrt{\frac{2\eta^3 x^3}{g\rho}} \sqrt{\frac{1}{t_g}} \left[\frac{1}{t_g} + \frac{1}{t_{rise}} \right]$$

Q: charge of oil drop

n: number of unpaired electrons in drop

e : elementary charge

d: plate separation

V: Voltage across plates

: density $\rho = \rho_{oil} - \rho_{air}$

 η : viscosity of air

g: gravitational constant

x: drift distance for oil drop

 t_g : fall time t_{rise} : rise time



Route of Charge Calculation Q(t_a, t_{rise}).

$$\frac{1}{f_c^{\frac{3}{2}}} \approx 1 - \left(\frac{t_g}{\tau_g}\right)^{\frac{1}{2}}; \quad \tau_g = \frac{2\eta x}{\rho g r_c^2}; \quad r_c[m] = \frac{6.18 \times 10^{-5}}{\rho [mmHg]}$$

$$Q = F \bullet S \bullet T = \left(\frac{1}{f_c^{3/2}}\right) \frac{9\pi d}{V} \sqrt{\frac{2\eta^3 x^3}{g\rho}} \frac{1}{\sqrt{t_g}} \left(\frac{1}{t_g} + \frac{1}{t_{rise}}\right)$$

$$F = \frac{1}{f_c^{3/2}} \approx 1 - \left(\frac{t_g}{\tau_g}\right)^{\frac{1}{2}} S = \frac{9\pi d}{V} \sqrt{\frac{2\eta^3 x^3}{g\rho}} T = \frac{1}{\sqrt{t_g}} \left(\frac{1}{t_g} + \frac{1}{t_{rise}}\right)$$

$$S = \frac{9\pi d}{V} \sqrt{\frac{2\eta^3 x^3}{g\rho}}$$

$$T = \frac{1}{\sqrt{t_g}} \left(\frac{1}{t_g} + \frac{1}{t_{rise}} \right)$$

Route of Charge Calculation. Origin Projects. Data Collecting.

Projects Section L1.opj ... Section L4.opg

Locations:

\\engr-file-03\PHYINST\APL Courses\PHYCS401\Common\Origin templates\Oil drop experiment \\engr-file-03\PHYINST\APL Courses\PHYCS401\Students\1.Millikan Oil Drop experiment

A(L)	B(Y)	C(Y)	D(Y)	E(Y)	F(Y)	G(Y)	H(Y)	
parameter label	Par	tg	tr	parameter label	Par	tg	tr	pa
student1, student2	student1, student2	student1, student2	student1, student2	student3, student4	student3, student4	student3, student4	student3, student4	5
η				η				η
Δη/ΔΤ				$\Delta\eta/\Delta T$				Δτ
ρ1				ρ1				ρ1
ρ2				ρ2				ρ2
ρ1–ρ2				ρ1–ρ2				ρ1
g								g
р								р
X				x				х
d				d				d
V				V				V
Та				Ta				Ta
								6
1 2 4	student1, student2 AA/AT p1 p2 p1-p2 g	student1, student2 An/AT p1 p2 p1-p2 p1-p2 p1-p2 p1-p2	student1, student1, student2 An/AT p1 p2 p1-p2 p1-p2 g px di	student1, student2 student1, student2 student2 An/AT p1 p2 p1-p2 g px did v/	Student1, Student1, Student1, Student2 Student2 Student2 Student2 Student4 Student4 Student4 Student2 Student4 Student4 Student5 Student6 Student6 Student7 Student7 Student8 Stude	Par tg tr parameter label Par student1, student2, student2, student2, student3, student4 π	Student1, Student1, Student2 Student2 Student3, Student4 Stude	Par tg tr parameter label Par tg tr parameter label Par tg tr


Route of Charge Calculation. Origin Projects. Data analysis.

Project: Millikan1.opj

\\engr-file-03.engr.illinois.edu\PHYINST\APL
Courses\PHYCS401\Common\Origin templates\Oil drop experiment


or

\\engr-file-03.engr.illinois.edu\PHYINST\APL
Courses\PHYCS401\Students\1. Millikan Oil Drop experiment






Route of Charge Calculation. Origin Project. Data Analysis.




Project Millikan1.opj

Plugin your data here




Route of Charge Calculation. Origin Project. Data Analysis.

Project Millikan1.opj



Route of Charge Calculation. Origin Project. Data Analysis.




Follow correct order of calculations: $r_c \rightarrow \tau_g \rightarrow (F,S,T) \rightarrow Q \rightarrow n$




Route of Charge Calculation. Origin Project. Data Analysis.


Project Millikan1.opj




Charge calculation. Origin project.



Expected results




Choice of Oil Drops for the Analysis: rise and fall times



Modern experiments at





- Drop generation rate 1 Hz
- Fluid Dow Corning silicon oil
- Number of drops 17 million
- Mass 70.1 milligrams
- Duration 8 months




Modern experiments at 5



Machine vision mediated auto-control of: the average charges of the drops, the fall path of the drops, the upward laminar air flow, and the electronic drive to the drop ejector are new features of this fluid drop charge measurement system.

The experiment is ran from 2004 - 2007.

Charge measurement accuracy achieved is better than 1/24 e for drops of up to 26 microns in diameter.




Modern experiments at




Summary as of January 2007.

Total mass throughput for all experiments- 351.4 milligrams of fluid Total drops measured in all experiments - 105.6 million

No evidence for fractional charge particles was found.





Appendix #1

Traditional reminder:

L1_Lab3_student name.pdf

Report-exp2.pdf

Please upload the files in proper folder! This week folders:

Pulses in transmission lines_L1

Pulses in transmission lines_L3

Pulses in transmission lines_L4


Pulses in transmission lines_L5

This week you have the last chance to submit "Transients in RLC" report



Appendix #2

Transmission line. Unknown load simulation




Location: \\Phyaplportal\PHYCS401\Common\Transmission line software



Appendix #2

Transmission line. Unknown load simulation



Location: \\Phyaplportal\PHYCS401\Common\Transmission line software

