


Fundamental Symmetry and Neutrino Physics

Liang Yang
Physics 403


April 12, 2022



Do we understand the Universe we live in?




Standard Cosmological Model




Standard Model of
Particle Physics

Standard Models are incomplete...

- What's the origin of matter – antimatter asymmetry in today's Universe?
- What is dark matter or dark energy?
- What is the nature of gravity?
- Can all forces in nature be unified?



Matter-Antimatter Asymmetry



Dark Matter

In Search of “New” Standard Model


■ LHC: direct search for new particles

- ◆ Discovery of Higgs!
- ◆ Hints of New Physics?

■ Precision measurements:

- ◆ EDMs of e , n , atoms, etc.
- ◆ Weak mixing angle
- ◆ $0\nu\beta\beta$
- ◆ Muon $g-2$
- ◆ Lepton flavor violation
- ◆ π , K and B decays
- ◆ Unitarity tests


Fermilab Muon g-2 result – new physics?



Theory: g-fac.: 2.00233183620(86) anomalous mag. moment: 0.00116591810(43)


Experiment: g-fac.: 2.00233184122(82), anomalous mag. moment: 0.00116592061(41)
5

Fermilab CDF new W mass Measurement




Seven standard deviations away from the theoretical prediction.


Neutrino Oscillation and Neutrino Mass




Super-K



SNO



K2K



KamLand

Super-K: atmospheric ν_μ neutrino oscillation

SNO: solar ν_e flavor transformation


K2K: accelerator ν_μ oscillation

Kamland: reactor $\bar{\nu}_e$ disappearance and oscillation


Neutrinos have Mass

The first evidence of physics beyond the Standard Model!

Our Sun is a copious source of electron type neutrinos ...




Experiment located 1500m underground
Homestake Gold Mine in SD
3 million times less cosmic ray interactions
(**bkgnds**) due to muons (which are very
penetrating particles), compared to the surface.



Far too few (~1/3) solar neutrinos were seen
compared to predicted solar production !

The plot thickens – some good fortune ...

1983 experiments (for protons decay) also good neutrino detectors ... cross check Homestake.




A massive detector, known as “SuperK”, clearly observed ν 's from the Sun, and confirmed the signal of missing solar ν 's.

In addition, SuperK was able to observe ν 's produced in the upper atmosphere by cosmic rays – “atmospheric ν 's”, and to tell where they were coming from, leading to a : **Breakthrough Observation in 1998**

In the Kamioka Mine in Japan

- Depth of 1000m
- Water tank (3000 tons for the first one)
- Instrumented to observe light flashes from produced from μ 's or e 's.
(led by M. Koshiba, also a 2002 Nobelist)



Particles are produced along the ν direction :
For the first time **directional information**.




(An aside : An unexpected dividend at Kamioka

The luckiest break since 1604 ! : Super Nova SN1987A
10⁵⁸ neutrinos produced from 168,000 light years away.
11 observed in 13 second interval by KamiokaNDE II)




Atmospheric Neutrino

Atmospheric neutrinos originate in cosmic ray “showers”



The **showers** produce electron, muon (and tau) type neutrinos, in a mix that can be predicted.

Neutrinos can reach SuperK from *above* or from *below* (the Earth is hardly a barrier at all to a neutrino, after all.)




Stunning Result


Observed neutrino rates & mixture differed for the two directions !

Were flavors changing in transit ?


SNO




2002 Sudbury Neutrino Observatory



Electron Scattering



Charge Current




Neutral Current

Unknown Properties of Neutrinos


Major Questions in Neutrino Physics

- Majorana particle, (i.e. its own antiparticle)
- Absolute mass scale of neutrinos.
- Mass hierarchy
- CP violation phase
- Anomalies (Sterile neutrinos?)




Double Beta Decay


Observable if single beta decay is forbidden



Two neutrino double beta decay




Neutrinoless double beta decay




Observation of $0\nu\beta\beta$:

- Majorana neutrino
- Neutrino mass scale
- Lepton number violation

The EXO-200 Detector




Liquid Xenon Time Projection Chamber




The EXO-200 time projection chamber uses both scintillation and ionization signals to fully reconstruct energy depositions inside liquid xenon

Single Site Events (SS)




Multiple Site Events (MS)



Event topology is a powerful tool not only for gamma background rejection, but also for signal discovery.

EXO-200 installation site: WIPP



- EXO-200 installed at WIPP (Waste Isolation Pilot Plant), in Carlsbad, NM
 - 1600 mwe flat overburden (2150 feet, 650 m)
 - U.S. DOE salt mine for low-level radioactive waste storage
 - Cleanroom installed on adjustable stands to compensate salt movements.
 - Salt “rock” low activity relative to hard-rock mine

$$\Phi_\mu \sim 1.5 \times 10^5 \text{ yr}^{-1} m^{-2} sr^{-1}$$


$$U \sim 0.048 \text{ ppm}$$

$Th \sim 0.25 ppm$

$$K \sim 480 \, ppm$$

Esch et al., arxiv:astro-ph/0408486 (2004)

EXO-200 $0\nu\beta\beta$ Results



EXO-200 TPC

Combined Phase I + II:

Limit $T_{1/2}^{0\nu\beta\beta} > 3.5 \times 10^{25}$ yr (90% C.L.)

$\langle m_{\beta\beta} \rangle < (93 - 286)$ meV

Sensitivity 5.0×10^{25} yr


2012: *Phys. Rev. Lett.* 109, 032505

2014: *Nature* 510, 229-234

2018: *Phys. Rev. Lett.* 120, 072701


2019: *Phys. Rev. Lett.* 123, 161802

- EXO-200 uses liquid xenon time projection chamber (TPC) to search for $0\nu\beta\beta$ of ^{136}Xe
- Successful operation from 2011 – 2018 with total ^{136}Xe isotope exposure of 234.1 kg·yr.
- Experimental sensitivities continue to exceed statistics due to improvements in hardware and analysis.
- Setting one of the strongest limits on this rare decay.




From EXO-200 to nEXO


EXO-200 as a technology demonstrator




2.5MeV γ
attenuation length
8.5cm = —



150kg





nEXO: a 5000 kg enriched LXe TPC




Pre-Conceptual Design of nEXO

- **5 tones of single phase LXe TPC.**
- Ionization charge collected by anode.
- 178nm lights detected by **~4 m² SiPM array** behind field shaping rings.
- Combining light and charge to enhance the energy resolution.


nEXO pre-CDR, arXiv:1805.11142




Tagging $\beta\beta$ decay daughter Ba




$\leq 58\text{-atom}$





$\leq 15\text{-atom}$




$\leq 4\text{-atom}$



0-atom



nEXO Sensitivity (with Ba tagging)



[nEXO Sensitivity Paper: arXiv:1710.05075](#)

Normal hierarchy

GCM: Rodriguez, Martinez-Pinedo,
Phys. Rev. Lett. 105 (2010) 252503

What can Neutrino tell us about the Universe?

- What role did neutrino play in the evolution of the universe? (~ 4% mass of the universe, absolute mass scale? Number of species? ... double beta decay experiment, tritium decay experiment, sterile neutrino search...)
- Can neutrino be responsible for the matter and anti-matter asymmetry? (CP violation phase? ... long baseline neutrino experiment)
- Neutrino might be the best probe deep into the universe (IceCube...)
- Supernovae neutrinos, relic neutrinos...