

## QED with quarks

We will now augment the QED Lagrangian with the remaining fermions,

$$\mathcal{L} \supset \sum_{f=1}^3 \bar{Q}_f \gamma^\mu D_\mu Q_f + u_R^{f\dagger} \sigma^\mu D_\mu u_R^f + d_R^{f\dagger} \sigma^\mu D_\mu d_R^f - Y_{ij}^d Q_i^\dagger H d_{Rj} - Y_{ij}^u Q_i^\dagger \hat{H} u_{Rj}$$

Just like in QED, where  $H \rightarrow \begin{pmatrix} 0 \\ v \end{pmatrix}$  and leptons got mass and electric charge, same thing happens for quarks:

$$Y_{ij}^d Q_i^\dagger H d_{Rj} \rightarrow m_{df} d_L^f d_R^f$$

$$Y_{ij}^u Q_i^\dagger \hat{H} u_{Rj} \rightarrow m_{uf} u_L^f u_R^f$$

Recall hypercharges:  $Y = \frac{1}{6}$  for  $Q$ ,  $Y = \frac{2}{3}$  for  $u_R$ ,  $Y = -\frac{1}{3}$  for  $d_R$

$$\text{Electric charge is } T_3 + Y = \begin{cases} \frac{1}{2} + \frac{1}{6} = \frac{2}{3}, u_L \\ -\frac{1}{2} + \frac{1}{6} = -\frac{1}{3}, d_L \\ 0 + \frac{2}{3} = \frac{2}{3}, u_R \\ 0 + (-\frac{1}{3}) = -\frac{1}{3}, d_R \end{cases}$$

$\Rightarrow$  in the Standard Model, up-type quarks are charge  $\frac{2}{3}$  fermions, down-type quarks are charge  $-\frac{1}{3}$ . We will describe experiments which test both spin and charge.

Note: quarks also interact with  $SU(3)_c$  gauge field. We will ignore this for now and pick it back up after the break.

$$\Rightarrow \mathcal{L}_{\text{quarks}} = \sum_{f=1}^3 \left( \bar{u}_f (i\cancel{\partial} + \frac{2}{3} eA) u_f + \bar{d}_f (i\cancel{\partial} - \frac{1}{3} eA) d_f - m_{uf} \bar{u}_f u_f - m_{df} \bar{d}_f d_f \right)$$


Only new Feynman rule is factor of  $\frac{2}{3}$  or  $\frac{1}{3}$  on quark-quark-photon vertex.

In the 1960's, it was hypothesized that the proton is a bound state of three quarks,  $p = uud$ . Let's see how to test this.

$$\text{Charge } +1 = +\frac{2}{3} + \frac{2}{3} - \frac{1}{3}$$

# Deep inelastic scattering

Consider the process  $e^- p \rightarrow e^- X$ , where  $X$  is any collection of final-state particles. We want to calculate the differential cross section in terms of only the electron's kinematic variables, so that we don't even have to observe  $X$ .



Strictly speaking, this is not a Feynman diagram in QED, which is why there is a blob at the proton-photon-X vertex. However, we can use the same tricks from last week to parameterize it.

$$\langle |M|^2 \rangle = e^2 L^{\mu\nu} W_{\mu\nu}, \text{ where (assuming } E_e \gg m_e \text{ so we can neglect } m_e \text{)}$$

$$L^{\mu\nu} = \frac{1}{2} \sum_{\text{spins}} \bar{u}(k') \gamma^\mu u(k) \bar{u}(k) \gamma^\nu u(k') = \frac{1}{2} \text{Tr}[\not{k}' \gamma^\mu \not{k} \gamma^\nu]$$

$$= 2(k'^\mu k^\nu + k'^\nu k^\mu - k \cdot k' \eta^{\mu\nu})$$

$$E_n E_V^\mu W^{\mu\nu} = \frac{1}{2} \sum_{X, \text{spins}} \int d\pi_X (2\pi)^4 \delta(q + P - P_X) |M(\gamma^\mu p \rightarrow X)|^2$$

avg. over proton spins

In general, we can't compute  $W$ , but it can only depend on  $P$  and  $q$ , and it must be symmetric and satisfy  $q_\mu W^{\mu\nu} = 0$ . There are two independent Lorentz scalars,  $q^2$  and  $P \cdot q$  ( $P^2 = m_p^2$  is a constant).

Conventional to take  $Q = \sqrt{-q^2}$  and  $x = \frac{Q^2}{2P \cdot q}$

$$\Rightarrow W^{\mu\nu} = W_1(Q, x) \times \left( -\eta^{\mu\nu} + \frac{q^\mu q^\nu}{q^2} \right) + W_2(Q, x) \times \left( P^\mu - \frac{P \cdot q}{q^2} q^\mu \right) \left( P^\nu - \frac{P \cdot q}{q^2} q^\nu \right)$$

with  $W_1$  and  $W_2$  unknown functions of the two independent variables  $Q$  and  $x$ .

Contract with  $L^{\mu\nu}$ , and specify to the lab frame where  $P = (m_p, 0, 0, 0)$ ,

$$k = (E, 0, 0, E), \quad k' = (E', E' \sin \theta, 0, E' \cos \theta).$$

$$\begin{aligned}
 (k'^\mu k^\nu + k'^\nu k^\mu - \eta^{\mu\nu} k \cdot k') (-\eta_{\mu\nu} + \frac{q_\mu q_\nu}{q^2}) &= -2k \cdot k' + 4k \cdot k' + \frac{2(q \cdot k)(q \cdot k')}{q^2} - k \cdot k' \\
 &= \frac{2((k-k') \cdot k)((k-k') \cdot k')}{(k'-k)^2} + k \cdot k' \\
 &= \frac{-2(k' \cdot k)^2}{-2k' \cdot k} + k' \cdot k \\
 &= 2k' \cdot k = 2EE'(1 - \cos \theta) \propto \sin^2 \frac{\theta}{2}
 \end{aligned}$$

Similarly, contracting the  $W_2$  term gives  $\cos^2 \frac{\theta}{2}$ .

What remains is  $\frac{d\sigma}{d\cos\theta dE'}$ , which is proportional to (putting in  $m_p$  for dimensions):

$$\frac{1}{m_p} W_1(Q, x) \sin^2 \frac{\theta}{2} + \frac{m_p}{2} W_2(Q, x) \cos^2 \frac{\theta}{2}.$$

$(Q, x) \leftrightarrow (E', \theta)$ , so by measuring the number of electrons scattered at energy  $E'$  and angle  $\theta$ , we can read off  $W_1$  and  $W_2$ .

However, experimentally it is found that  $W_1$  depends only on  $x$ , not  $Q$ .

We can understand this as a consequence of the proton "containing" point-like spin- $\frac{1}{2}$  particles, which we identify as quarks.

First, let's examine the elastic cross section  $e^- q_i \rightarrow e^- q_i$  where  $q_i$  has electric charge  $Q_i e$ . You will do this in HW:

$$\frac{d\sigma}{d\Omega} = \frac{e^4 Q_i^2}{64\pi^2 E^2 \sin^4 \frac{\theta}{2}} \frac{E'}{E} \left( \cos^2 \frac{\theta}{2} + \frac{E'-E}{m_{q_i}} \sin^2 \frac{\theta}{2} \right) \text{ for } E \gg m_e,$$

in lab frame where the quark is initially at rest.

For elastic scattering,  $p_q + q = p'_q$  where  $p_q$  and  $p'_q$  are quark initial/final momenta.

Squaring,  $m_{q_i}^2 - Q^2 + 2p_q \cdot q = m_{q_i}^2$ , so  $Q^2 = 2p_q \cdot q = 2m_{q_i}(E-E')$ .

This means we can write  $\frac{d\sigma}{d\Omega dE'} = \frac{\alpha^2 Q_i^2}{4E^2 \sin^4 \frac{\theta}{2}} \left[ \cos^2 \frac{\theta}{2} + \frac{Q^2}{2m_{q_i}} \sin^2 \frac{\theta}{2} \right] \delta\left(E-E'-\frac{Q^2}{2m_{q_i}}\right)$

(you will verify this in HW)

We will now make two assumptions:

19

$$1. \sigma(e^-p \rightarrow e^-X) = \sum_i \int_0^1 d\xi f_i(\xi) \hat{\sigma}(e^-q_i \rightarrow e^-q_i) \quad \text{with } p_{q_i} = \xi P$$

Proton is composed of quarks  $q_i$ , each of which has a random fraction of the proton's momentum  $\xi$  given by its parton distribution function  $f_i(\xi)$

2. Quarks are weakly interacting inside the proton, at large  $Q$ .

This seems weird: isn't the strong force "strong"? How would quarks bind together to make the proton if this were true?

More on this after the break!

From  $p_{q_i} = \xi P$ , and  $P = (m_p, 0, 0, 0)$ , we set  $m_{q_i} = \xi m_p$  in previous formulas.  $P \cdot q = m_p(E - E')$ , so  $x = \frac{Q^2}{2P \cdot q} = \frac{Q^2}{2m_p(E - E')}$ , and

$$\delta(E - E' - \frac{Q^2}{2m_{q_i}}) = \delta\left(\frac{Q^2}{2m_p x} - \frac{Q^2}{2m_p \xi}\right) = \frac{2m_p}{Q^2} \delta\left(\frac{1}{x} - \frac{1}{\xi}\right) = \frac{2m_p x^2}{Q^2} \delta(\xi - x)$$

$$\Rightarrow \frac{d\sigma(e^-p \rightarrow e^-X)}{d\Omega dE'} = \sum_i f_i(x) \frac{\alpha^2 Q_i^2}{4E^2 \sin^2 \frac{\theta}{2}} \left[ \frac{2m_p}{Q^2} x^2 \cos^2 \frac{\theta}{2} + \frac{1}{m_p} \sin^2 \frac{\theta}{2} \right]$$

$\Rightarrow W_2 \propto \sum_i Q_i^2 f_i(x)$ , only depends on  $x$ ! Proton looks the same no matter how hard it is hit.

This prediction is beautifully backed up by data. The interpretation is that the proton has point-like constituents. The ratio  $\frac{W_2}{W_1} = \frac{4x^2}{Q^2}$  is characteristic of spin- $\frac{1}{2}$  constituents, which is also confirmed by data.

Using deep inelastic scattering, we can measure  $F_2$  and  $F_L$ .

Momentum conservation should imply  $\sum_j \int_0^1 d\xi \xi f_j(\xi) = 1$ . However,

the measured value is 0.38! Most of the proton's momentum is carried by gluons - QCD is complicated!