

Quantum electrodynamics

17

SM Lagrangian from last time:

$$\begin{aligned} \mathcal{L}_{SM} &= \mathcal{L}_{kinetic} + \mathcal{L}_{Yukawa} + \mathcal{L}_{Higgs} \\ &= |D_\mu H|^2 - \frac{1}{4} G_{\mu\nu}^a G^{\mu\nu a} - \frac{1}{4} W_{\mu\nu}^a W^{\mu\nu a} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} \\ &\quad + \sum_{f=1}^3 \left\{ \underline{i L_f^\dagger \bar{\sigma}^\mu D_\mu L_f} + i Q_f^\dagger \bar{\sigma}^\mu D_\mu Q_f + \underline{i e_R^\dagger \sigma^\mu D_\mu e_R^f} + i u_R^\dagger \sigma^\mu D_\mu u_R^f + i d_R^\dagger \sigma^\mu D_\mu d_R^f \right\} \\ &\quad - \underline{Y_{ij}^e L_i^\dagger H e_R^j} - Y_{ij}^d Q_i^\dagger H d_R^j - Y_{ij}^u Q_i^\dagger \tilde{H} u_R^j + \text{h.c.} \\ &\quad + m^2 H^\dagger H - \lambda (H^\dagger H)^2 \end{aligned}$$

Focus on these terms today. After setting $H = \begin{pmatrix} 0 \\ v \end{pmatrix}$ and diagonalizing

Y_{ij}^e , bottom component of fermion doublet $L_f = \begin{pmatrix} \nu_f^f \\ e_f^f \end{pmatrix}$ is

$$\sum_{f=1}^3 i e_L^{f\dagger} \bar{\sigma}^\mu D_\mu e_L^f + i e_R^{f\dagger} \sigma^\mu D_\mu e_R^f - y_{f\nu} e_L^{f\dagger} e_R^f + \text{h.c.}$$

We want to identify $y_{f\nu} \equiv m_f$, but for this to describe charged leptons (electrons, muons, taus), we have to be able to combine L and R spinors into a 4-component spinor $\Psi = \begin{pmatrix} e_L \\ e_R \end{pmatrix}$ with the correct electric charge. Recall $Y = -1$ for e_R , but $Y = -\frac{1}{2}$ for e_L , so this isn't quite right.

In fact, $Q = T_3 + Y$, where T_3 is the 3rd generator of $SU(2)_L$

$T_3 = \frac{1}{2} \sigma_3 = \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix}$, so e_L is an eigenvector of T_3 w/ eigenvalue $-\frac{1}{2}$.

$$Q_L = -\frac{1}{2} + (-\frac{1}{2}) = -1 \quad \left. \vphantom{Q_L} \right\} \text{this works!}$$

$$Q_R = 0 + -1 = -1$$

Conclusion: electromagnetism is a (linear combination of $SU(2)_L$ and $U(1)_Y$ gauge bosons.

We will see later on that the remaining $SU(2)$ gauge fields are much heavier than m_e, m_μ , so for the time being we can ignore them.

$$\mathcal{L}_{QED} = \left\{ \sum_{f=1}^3 \bar{\Psi}_f (i \partial_\mu - e A_\mu) \gamma^\mu \Psi_f - m_f \bar{\Psi}_f \Psi_f \right\} - \frac{i}{4} F_{\mu\nu} F^{\mu\nu}$$

where $\Psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}$, $\bar{\Psi} = (\psi_R^\dagger \ \psi_L^\dagger) = \Psi^\dagger \gamma^0$

Classical spinor solutions

(Massive) Dirac Equation: $i \gamma^\mu \partial_\mu \Psi - m \Psi = 0$

Look for solutions $\Psi = e^{-i p \cdot x} \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix}$ where χ_L, χ_R are constant 2-comp spinors

$\Rightarrow \gamma^\mu p_\mu \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix} = m \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix}$

$$\begin{pmatrix} 0 & p \cdot \sigma \\ p \cdot \bar{\sigma} & 0 \end{pmatrix} \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix} = m \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix}$$

First look for solutions with $\vec{p} = 0$; can construct the solution for general \vec{p} with a Lorentz boost. $p \cdot \sigma = p \cdot \bar{\sigma} = m \mathbb{1}$, so

$$\begin{pmatrix} -\mathbb{1} & \mathbb{1} \\ \mathbb{1} & -\mathbb{1} \end{pmatrix} \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix} = 0 \quad \Rightarrow \chi_L = \chi_R, \text{ but otherwise unconstrained}$$

Choose a basis: $\chi_L = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ or $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, so let 4-component solutions be

$u_\uparrow = \sqrt{m} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$ and $u_\downarrow = \sqrt{m} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}$. These represent spin-up and spin-down electrons (or muons or taus)

Just like with complex scalar fields, there are also negative-frequency solutions $e^{+i p \cdot x} \begin{pmatrix} \chi_L \\ \chi_R \end{pmatrix}$ that represent antiparticles: positrons. Changing sign of p^0 means $\chi_L = -\chi_R$.

Note: different labeling convention from Schwartz.

$v_\uparrow = \sqrt{m} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}$, $v_\downarrow = \sqrt{m} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix}$

Physical spin-up positrons have $\chi_L = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$; this comes from QFT.

Can construct solution for general p with Lorentz transformations.

For now, will just write down the solution and check that it works:

$$u^s(p) = \begin{pmatrix} \sqrt{p \cdot \sigma} \xi_s \\ \sqrt{p \cdot \bar{\sigma}} \xi_s \end{pmatrix}, \quad v^s(p) = \begin{pmatrix} \sqrt{p \cdot \sigma} \eta_s \\ -\sqrt{p \cdot \bar{\sigma}} \eta_s \end{pmatrix} \quad \text{where } \xi_1 = \eta_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \xi_2 = \eta_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (s=1, 2)$$

Check Dirac equation for u :

$$\begin{pmatrix} 0 & p \cdot \sigma \\ p \cdot \bar{\sigma} & 0 \end{pmatrix} \begin{pmatrix} \sqrt{p \cdot \sigma} \xi_s \\ \sqrt{p \cdot \bar{\sigma}} \xi_s \end{pmatrix} = \begin{pmatrix} \sqrt{p \cdot \sigma} \sqrt{(p \cdot \sigma)(p \cdot \bar{\sigma})} \xi_s \\ \sqrt{p \cdot \bar{\sigma}} \sqrt{(p \cdot \sigma)(p \cdot \bar{\sigma})} \xi_s \end{pmatrix} = \begin{pmatrix} \sqrt{p \cdot \sigma} \sqrt{m^2} \xi_s \\ \sqrt{p \cdot \bar{\sigma}} \sqrt{m^2} \xi_s \end{pmatrix} = m u \sqrt{\quad}$$

useful identity

To see how the spinors behave, let's let $\vec{p} = p_z \hat{z}$:

$p \cdot \sigma = \begin{pmatrix} E - p_z & 0 \\ 0 & E + p_z \end{pmatrix}$, $p \cdot \bar{\sigma} = \begin{pmatrix} E + p_z & 0 \\ 0 & E - p_z \end{pmatrix}$, and since these matrices are already diagonal, taking the square root is unambiguous

$$u_1 = \begin{pmatrix} \sqrt{E - p_z} \\ 0 \\ \sqrt{E + p_z} \\ 0 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 0 \\ \sqrt{E + p_z} \\ 0 \\ \sqrt{E - p_z} \end{pmatrix}, \quad v_1 = \begin{pmatrix} \sqrt{E - p_z} \\ 0 \\ -\sqrt{E + p_z} \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 0 \\ \sqrt{E + p_z} \\ 0 \\ -\sqrt{E - p_z} \end{pmatrix}$$

* NOTE: very bad typo in Schwartz 2nd edition eq. (11.26)!

If $E \gg m$, $E \approx |p_z|$. For $p_z > 0$ (motion along $+z$ -axis),

$$u_1(p) \approx \sqrt{2E} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}. \quad x_L = 0, \text{ so this is a purely right-handed spinor}$$

But $\xi = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ means spin-up along z -axis; this electron also has helicity $+\frac{1}{2}$, or has right-handed polarization in the traditional sense.

\Rightarrow for massless particles, chirality and helicity are the same
(right-handed spinor = right-handed particle)

What about antiparticles? A positron moving in the +z direction with spin-up along z-axis is still a right-handed antiparticle, but its spin is

$$v_2(p) = \begin{pmatrix} 0 \\ \sqrt{E+p_z} \\ 0 \\ \sqrt{E-p_z} \end{pmatrix} \approx \sqrt{2E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \text{ which is pure } \chi_L. \text{ Helicity and chirality}$$

are opposite for antiparticles.

Think of u's and v's as column vectors and $\bar{u} \equiv u^\dagger \gamma^0$, $\bar{v} \equiv v^\dagger \gamma^0$ as row vectors.

Useful identities for what follows:

$$\begin{aligned} \bar{u}_s(p) u_{s'}(p) &= u_s^\dagger(p) \gamma^0 u_{s'}(p) = \begin{pmatrix} \xi_s^\dagger \sqrt{p \cdot \sigma} & \xi_s^\dagger \sqrt{p \cdot \bar{\sigma}} \end{pmatrix} \begin{pmatrix} \sqrt{p \cdot \bar{\sigma}} \xi_{s'} \\ \sqrt{p \cdot \sigma} \xi_{s'} \end{pmatrix} \\ &= \begin{pmatrix} \xi_s^\dagger & \xi_s^\dagger \end{pmatrix} \begin{pmatrix} \sqrt{p \cdot \sigma} \sqrt{p \cdot \bar{\sigma}} & \\ & \sqrt{p \cdot \bar{\sigma}} \sqrt{p \cdot \sigma} \end{pmatrix} \begin{pmatrix} \xi_{s'} \\ \xi_{s'} \end{pmatrix} = 2m \delta_{ss'} \end{aligned}$$

$$\text{Similarly, } u_s^\dagger(p) u_{s'}(p) = \begin{pmatrix} \xi_s^\dagger & \xi_s^\dagger \end{pmatrix} \begin{pmatrix} p \cdot \sigma & \\ & p \cdot \bar{\sigma} \end{pmatrix} \begin{pmatrix} \xi_{s'} \\ \xi_{s'} \end{pmatrix} = 2E \delta_{ss'} \text{ (note: not Lorentz-invariant!)}$$

Analogous for v (check yourself):

$$\bar{v}_s(p) v_{s'}(p) = -2m \delta_{ss'}, \quad v_s^\dagger(p) v_{s'}(p) = 2E \delta_{ss'}$$

We've been a bit fast and loose with matrix notation. The above were inner products: contract two 4-component spinors to get a number.

Can also take outer products to get a 4x4 matrix:

$$\sum_{s=1}^2 u_s(p) \bar{u}_s(p) = \not{p} + m \mathbb{1}_{4 \times 4} \equiv \not{p} + m \text{ (Feynman slash notation)}$$

$$\sum_{s=1}^2 v_s(p) \bar{v}_s(p) = \not{p} - m \quad \star \text{ HW} \quad \text{note the order of u and } \bar{u}, \text{ and same spin index!}$$

Classical vector solutions

Gauge-fixed Maxwell equations: $\square A_\mu = 0, \partial^\mu A_\mu = 0$

Again, look for solutions $A_\mu = \epsilon_\mu(p) e^{-ip \cdot x}$. We did this in week 4: in a frame where $p^\mu = (E, 0, 0, E)$, we have

$$\epsilon_\mu^{(1)} = (0, 1, 0, 0), \epsilon_\mu^{(2)} = (0, 0, 1, 0), \epsilon_\mu^f = (1, 0, 0, 1)$$

Recall ϵ_μ^f is unphysical because it has zero norm. However, we need to include it because $\epsilon_\mu^{(1,2)}$ mix with it under a Lorentz transformation.

Explicitly, let $\Lambda_\nu^\mu = \begin{pmatrix} \frac{3}{2} & 1 & 0 & -\frac{1}{2} \\ 1 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ \frac{1}{2} & 1 & 0 & \frac{1}{2} \end{pmatrix}$. Can check $\eta^\alpha \Lambda_\alpha^\mu \Lambda_\nu^\mu = \eta^{\mu\nu}$, also $\Lambda_\mu^\nu p^\nu = p^\mu$,

so Λ preserves p^μ . However, $\Lambda_\nu^\mu \epsilon_\mu^{(1)} = (1, 1, 0, 1) = \epsilon_\nu^{(1)} + \epsilon_\nu^f$, so Lorentz transformations can generate the unphysical polarization.

But it turns out that in QED, all amplitudes $M^\mu(p)$ involving an external photon with momentum p^μ satisfy $\boxed{p_\mu M^\mu = 0}$. This is the Ward identity, and because $\epsilon_\mu^f \propto p^\mu$, this unphysical polarization doesn't contribute to any observable quantity. (More on this later!)

Analogous to spinors, we can compute inner and outer products:

$$\epsilon_\mu^{(i)} \epsilon^{\mu(j)} = -\delta^{ij}, \quad i = 1, 2$$

$$\begin{aligned} \sum_{i=1}^2 \epsilon^{\mu(i)} \epsilon^{\nu(i)} &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = -\eta^{\mu\nu} + \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \\ &= -\eta^{\mu\nu} + \frac{p^\mu \bar{p}^\nu + p^\nu \bar{p}^\mu}{p \cdot \bar{p}} \end{aligned}$$

where $\bar{p} = (E, 0, 0, -E)$. But by the arguments above, the p^μ will always contract to zero, so we can say

$$\sum_{i=1}^2 \epsilon^{\mu(i)} \epsilon^{\nu(i)} \rightarrow -\eta^{\mu\nu} \quad (\text{again, sum over spins gives a matrix})$$