


## The $\nu = 5/2$ fractional quantum Hall effect

As we have seen, the vast majority of quantum Hall plateaux observed experimentally, whether in Si MOSFET's, GaAlAs-GaAs heterostructures or graphene, occur either at integral values of the filling factor  $\nu \equiv N_e/N_\phi$  ( $N_e \equiv$  number of electrons,  $N_\phi \equiv$  number of flux quanta) (IQHE) or at rational fractions  $\nu = p/q$  with  $q$  an *odd* integer (FQHE). Within the composite-fermion picture, the FQHE is explained by setting  $p = k$ ,  $q = 2nk \pm 1$  and viewing the FQHE as derived from the IQHE with  $\nu = k$  by attaching to each electron  $2n$  imaginary “flux quanta”; or, what is equivalent, multiplying the IQHE wave function by  $(z_i - z_j)^{2n}$ . However, as we have seen in lecture 20, in the case of  $\nu = 1/2$  the same procedure yields the conclusion that the “parent” IQHE state should correspond to  $\nu = \infty$ , i.e. it should be a Fermi liquid. Moreover, we saw that experiments, in particular on magnetic focusing, seem consistent with this point of view.

Prima facie, one would expect the states which occur at  $\nu = n + 1/2$  in traditional QHE systems such as GaAlAs-GaAs heterostructures<sup>1</sup> to differ from that at  $\nu = 1/2$  only by having  $n$  Landau levels completely filled; the situation in the  $(n + 1)$ -th LL should be similar to that in the LLL for  $\nu = 1/2$ . It was therefore a considerable surprise when it was discovered in 1987 that a QH plateau occurs, with all the standard characteristics, at  $\nu = 5/2$ . This plateau seems to be quite robust, with a gap  $\Delta \sim 500$  mK; at temperatures  $\ll \Delta$  the longitudinal resistance vanishes within experimental accuracy, and the Hall conductance appears to be quantized at  $(5/2)e^2/h$  to high accuracy, thereby excluding the possibility that the plateau is a case of the standard FQHE with  $\nu = 32/13$  or  $33/13$ . To date, the only other even-denominator plateau which has been reliably seen is at  $\nu = 7/2$ , which might naturally be viewed as similar to that at  $\nu = 5/2$  with the  $n = 2, \uparrow$  spin LL filled.  $\nu = 19/8$  gives a deep dip in  $R_{xx}$ , and a barely visible plateau. (In 1-st LL, not only  $\nu = 1/2$  but also  $\nu = 1/4, 3/4$  give apparently Fermi-liquid-like states).

Some other properties of the  $\nu = 5/2$  state:

1. Robustness: both plateau in  $R_H$  and the zero of  $R_{xx}$  extend over a range  $\sim 0.1$  in  $\nu$ . (in the highest-mobility samples)<sup>2</sup>
2. Excitation gap  $\Delta$ : this is measured by fitting  $R_{xx}$  to  $R_{xx} \sim \text{const. exp } -\Delta/T$ .  $\Delta$  appears to be a strong ( $\sim$  exponential) function of disorder: the highest measured value to date  $\sim 0.45$  K, and extrapolation to zero disorder ( $\mu \rightarrow \infty$ ) gives  $\Delta \sim 0.6$  K ( $\sim 0.006V_c$ ) ( $V_c = e^2/4\pi\epsilon\epsilon_0l_M$ ) (Note:  $V_c$  is the natural unit to measure qp gap, as  $\Delta_{qp} \equiv 0$  for the FQHE if  $V_c$  is neglected).
3. Magnetic field dependence:<sup>3</sup> If  $B_\perp$  (hence  $\nu$ ) is held constant, and  $B_\parallel$  is varied,  $\Delta_{qp}$



<sup>1</sup>As we saw in lecture 22, a QH plateau would occur at  $\nu = n + 1/2$  in (single-layer) graphene if both the spin and valley degeneracies were split, but this has to do with the special nature of the Dirac spectrum and is best regarded as a variant of the IQHE.

<sup>2</sup>See fig. 1 of Xia et al., PRL **93**, 176809 (2004).

decreases linearly with  $B_{\parallel}$ , extrapolating to 0 at  $B_{\parallel} = 1.5 - 2.5$  T. In the same geometry,  $\nu = 7/2$  FQHE shows similar behavior, but for  $\nu = 7/3$  (“Laughlin” state)  $\Delta_{\text{qp}}$  *increases* with  $B_{\parallel}$ .

4. At least to date, the only system in which the  $\nu = 5/2$  QH plateau has been seen is GaAs heterostructures: here typical parameters are  $n_s \sim 1 - 3 \times 10^{11} \text{ cm}^{-2}$  (so  $B \sim 2 - 6$  T),  $\mu \sim 3 \times 10^7 \text{ cm}^2/\text{V sec}$ ,  $T \sim 5 - 100$  mK.

In trying to understand what is going on at  $\nu = 5/2$  it seems very natural to make the default assumption that the LLL is filled both for  $\uparrow$  and  $\downarrow$  spin states; then prima facie the behavior in the  $n = 1, \uparrow$  spin LL should be identical to that in the  $n = 0, \uparrow$  spin LL for  $\nu = 1/2$ . However, this need not necessarily be the case, because in view of the different behavior of the wave functions for  $n = 0$  and  $n = 1$  the relevant matrix elements of the Coulomb interaction could be appreciably different in the two LL’s (note by the way that no FQHE has (to date) ever been seen for  $\nu > 4$ , i.e. presumably, in the third LL). An immediate question which arises then is: given that the LLL is completely filled for both spins, i.e. unpolarized, is the  $n = 1$  LL spin-polarized or not? As we shall see below, the answer to this question is crucial to the identification of the nature of the wave function and thus to the possibility of using the  $\nu = 5/2$  FQHE for TQC; unfortunately, it has not so far proved possible to determine the answer experimentally. Originally, it was found that the plateau is suppressed by a substantial magnetic field component parallel to the plane, and the most obvious explanation is that the relevant state is a spin singlet (hence energetically disadvantaged by the Zeeman field). However, subsequent experiments, combined with numerical theory, tend to suggest that the effect is actually of orbital origin; because the parallel component of the magnetic field affects the motion *perpendicular* to the plane, it can change the relevant Coulomb matrix elements. This raises the very obvious question: Does a corresponding QH effect occur in graphene? Assuming that at high fields the spin degeneracy is split but the valley degeneracy remains unsplit, the value of  $\nu$  corresponding to the  $5/2$  in the standard systems, i.e. the value at which the second LL with spin  $\uparrow$  is half filled, would be  $\nu = (\pm)3$ . There is in fact evidence for a  $\nu = 3$  FQHE in very recent experiments,<sup>4</sup> but it has not been so far investigated in detail and could be due to valley splitting. For the moment, let us assume that the  $\nu = 5/2$  state seen in GaAs heterostructures is in fact fully spin polarized and ask what is its nature? In particular, why does a QH system at  $\nu = 5/2$  not behave, as it seems to at  $\nu = 1/2$ , as a “disguised” Fermi liquid (of composite fermions)?

A relevant question is: What do we know about the possible instabilities of a Fermi liquid? There are of course a great many, but most of them, such as crystallization, tend to occur either not at all or at temperatures comparable to the Fermi temperature, which for GaAs heterostructures at  $n = 10^{11} \text{ cm}^{-2}$  is  $\sim 30$  K. The obvious instability which occurs for arbitrary weak interactions of the right sign and thus at arbitrary low temperatures is *Cooper pairing*, which of course in a system of real particles leads to superconductivity (if charged) or superfluidity (if neutral). This consideration led

<sup>3</sup>Dean et al., PRL **101** 186806 (2008).

<sup>4</sup>The experiments of the Columbia and Rutgers groups cited in lecture 22.

Moore and Read<sup>5</sup> to conjecture that

the  $\nu = 5/2$  QH plateau corresponds to a Cooper-paired state of composite fermions.

If this is true, then it is generally believed that the elementary excitations will be non-abelian (Ising) anyons, which is what makes this possibility so interesting in the context of TQC.

Let's try to make this hypothesis a bit more quantitative. According to the composite-fermion hypothesis, the correct groundstate for a given value of  $\nu = k/(2nk \pm 1)$  is given by taking the non-Gaussian part of the wave function of electrons at  $\nu = k$ , multiplying by  $(z_i - z_j)^{2n}$  ("adding  $2n$  flux quanta to each electron") and readjusting the Gaussian part so that the magnetic length  $l_M$  which comes in refers to the actual magnetic field  $B$  (not  $B^*$ ). Now, consider the case  $\nu = 1/2$ . This corresponds to the choice  $k \rightarrow \infty$ ,  $n = 1$ ; since in this limit  $B^* = 0$ , the wave function of a set of *noninteracting* electrons is just the filled Fermi sea for given value of  $n_s$  and complete spin polarization |FS>. Consequently, according to the above prescription we should have apart from normalization

$$\Psi_{\nu=1/2}^{(\text{normal})} = \prod_{ij} (z_i - z_j)^2 \exp - \sum_i |z_i|^2 / 4l_M^2 \text{ |FS} \rangle \quad (1)$$

where the |FS> component guarantees the correct antisymmetry under the exchange  $i \leftrightarrow j$ . The assumption which seems to be implicit in much of the theoretical literature is that a similar expression, multiplied by the appropriate Slater determinants for the completely filled (spin singlet) LLL wave function, would be adequate also for the  $\nu = 5/2$  state were it not for the effect of interactions. In the following I will not write out the part of the wave function which refers to the LLL explicitly.

So, if eqn. (1) is the correct representation of a normal Fermi sea of composite fermions, what is the corresponding representation for a BCS-paired state? The obvious answer is

$$\Psi_{\nu=1/2}^{(\text{paired})} = \prod_{ij} (z_i - z_j)^2 \exp - \sum_i |z_i|^2 / 4l_M^2 \text{ |BCS} \rangle \quad (2)$$

where |BCS> is the Cooper-paired state of weakly interacting fermions at the relevant density. Now, on our assumption that the  $\nu = 5/2$  state (as well as the  $\nu = 1/2$  state) is completely spin-polarized, the Fermi antisymmetry requires that the pairing takes place in a state of odd relative orbital angular momentum  $l$ , and the default option is  $l = 1$  ( $p$ -state). Moreover, the state must be two-dimensional.<sup>6</sup> This still does not specify the state uniquely; for example, the order parameter could be of the form  $Ak_x$  (or  $Ak_y$ ), which breaks rotational invariance but not time-reversal invariance. However, our general experience with BCS pairing suggests that in a rotationally invariant system it is usually energetically advantageous to make the energy gap (which is proportional

<sup>5</sup>Nuc. Phys. **360**, 362 (1991).

<sup>6</sup>Of course, as we have seen in lecture 9, no true superfluid ODLRO can survive in 2D. However, we may assume, at least for the moment, that we are below the KT transition, so that the lack of ODLRO does not affect the qualitative behavior.

to the modulus of the order parameter) as uniform as possible over the Fermi surface. This suggests that we should choose for the OP

$$\Delta(\mathbf{k}) = \Delta_0(k_x \pm ik_y) \quad (3)$$

so that even though the magnetic field  $B^*$  acting on the composite fermions is zero, their state still breaks time reversal invariance. In the literature a state of the form (3) is referred to as a “ $p + ip$ ” state: note that the “energy gap” is independent of  $\mathbf{k}$  and equal to  $\Delta_0$ .

The crucial question, now, is: What is the explicit form of the groundstate wave function  $|\text{BCS}\rangle$  which corresponds to the choice (3)? Actually, as we shall see in the next lecture, the answer does not seem to be unique in the thermodynamic limit (a fact which has not been widely appreciated in the QH literature). However, there is a particular answer which has been widely given in the literature on superfluid  $^3\text{He}$  and other (non-FQH) condensed matter systems, namely that the wave function is of the standard BCS form, with the coefficients  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  having the right angular dependence for a  $p + ip$  state. This is, explicitly,

$$|\text{BCS}\rangle = |\text{BCS}\rangle_{\text{standard}} \equiv \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{-\mathbf{k}}^\dagger) |0\rangle \quad (4)$$

where  $|0\rangle$  is the physical vacuum, the spin suffix  $\uparrow$  is omitted for simplicity, and the coefficients  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  are given by

$$u_{\mathbf{k}} = \frac{1}{\sqrt{2}}(1 + \epsilon_{\mathbf{k}}/E_{\mathbf{k}}), \quad v_{\mathbf{k}} = \frac{1}{\sqrt{2}}(1 - \epsilon_{\mathbf{k}}/E_{\mathbf{k}}) \exp i\phi_{\mathbf{k}} \quad (5)$$

$$E_{\mathbf{k}} \equiv \sqrt{(\epsilon_{\mathbf{k}} - \mu)^2 + \Delta_0^2}$$

with  $\mu$  the chemical potential. It is easily verified that the expectation value of the total orbital angular momentum  $L$  in the state described by  $|\text{BCS}\rangle_{\text{standard}}$  is  $N\hbar/2$ .

Should we simply insert  $|\text{BCS}\rangle_{\text{standard}}$  in the conjecture (2) for the groundstate of the  $\nu = 5/2$  QH system? There is a problem here, since we do not know a priori the value of the effective Coulomb matrix elements for the composite-fermion states and hence cannot calculate the gap magnitude  $\Delta_0$ . However, since the order of magnitude of the Coulomb energy ( $e^2/4\pi\epsilon_0 r_0$ , where  $r_0 \sim n_s^{-1/2}$ ) is quite comparable to the Fermi energy and may in fact be larger, it seems reasonable to suppose that  $\Delta_0$  is of order  $\epsilon_F$  and thus that the pair radius  $\xi$  of the pairs in the state  $|\text{BCS}\rangle_{\text{standard}}$  is of order of the interparticle spacing (or the magnetic length, which for  $\nu \sim 1$  is essentially the same thing). But there is little reason to believe that the composite-fermion idea works quantitatively on this kind of scale. Consequently, it may seem sensible to insert in (2) not the full real-space wave function derived from (4), but only the form which the latter takes at long distances ( $|\mathbf{r}_i - \mathbf{r}_j| \gg k_F^{-1}, \xi$ ). As we shall see in the next lecture, this has the form of the “Pfaffian”

$$\text{Pf} \left( \frac{1}{z_i - z_j} \right) \equiv \frac{1}{z_1 - z_2} \frac{1}{z_3 - z_4} \frac{1}{z_5 - z_6} \dots - \frac{1}{z_1 - z_3} \frac{1}{z_2 - z_4} \frac{1}{z_5 - z_6} \dots + \dots - \dots \quad (6)$$

i.e. it is the completely antisymmetrized version of the expression  $\prod_{j=i+1}^N \left(\frac{1}{z_i - z_j}\right)$ .

Thus, finally, the ansatz of Moore and Read for the  $\nu = 5/2$  QH state is up to normalization (presumably omitting the filled LLL)

$$\Psi_{\text{MR}}\{z_i\} = \prod_{i<j} (z_i - z_j)^2 \text{Pf} \frac{1}{(z_i - z_j)} \exp - \sum_i |z_i|^2 / 4l_M^2 \quad (7)$$

and it is on this conjectured form that most of the work on the possible implementation of TQC in this system has been based. Note that apart from the Pfaffian factor,  $\Psi_{\text{MR}}$  is just the Laughlin factor for  $\nu = 1/2$ ; however, the Pfaffian factor is of course essential to give the correct antisymmetry.

Although the ansatz (7) is an informal guess, it can be made plausible by two considerations:

1. It is the exact groundstate of the artificial Hamiltonian

$$\hat{H} = +V_0 \sum_{ijk} \delta^{(2)}(\mathbf{r}_i - \mathbf{r}_j) \delta^{(2)}(\mathbf{r}_j - \mathbf{r}_k) \quad (\delta^{(2)}(\mathbf{r}) \equiv \delta(x)\delta(y)) \quad (8)$$

2. Numerical studies show that it has a substantial overlap with the exact groundstate of some rather more realistic model Hamiltonians.

It is sometimes said in the literature that the MR state (7) is the exact analog of the superfluid  $A_1$  state of liquid  $^3\text{He}$ . This is not strictly true, since as well as the paired up-spin component  $^3\text{He-}A_1$ , also has an unpaired down-spin component. It would be true for (hypothetical) totally spin-polarized  $^3\text{He-}A_1$ , with no down-spin component. Note by the way that while the  $\nu = 1/2$  state would appear to be symmetric with respect to particles and holes, the Hamiltonian (8) breaks the particle-hole symmetry, and since the MR state is an exact eigenfunction of it, it must do the same.

Let's briefly review some other possible identifications of the  $\nu = 5/2$  QH state.<sup>7</sup>

1. The (331) state

Like the MR state, this is a triplet-paired state of the composite fermions; however, unlike that state (which corresponds to  $S = 1, S_z = 1$ ) this one corresponds to  $S = 1, S_z = 0$  and hence has no net spin polarization in any direction. The explicit form of this state in terms of the electron coordinates is

$$\Psi = \sum_{\{\sigma_i\}: \sum_i \sigma_i = 0} \prod_{\substack{i<j \\ \sigma_i = \sigma_j}} (z_i - z_j)^3 \prod_{\substack{i<j \\ \sigma_i \neq \sigma_j}} (z_i - z_j)^1 \exp - \sum_i |z_i|^2 / 4l_M^2 \quad (9)$$

In words: the correlation of any two parallel-spin electrons vanishes as  $r^3$ , but correlation of any two anti-parallel spin electrons vanishes only as  $r$ .

<sup>7</sup>The most readable account of this subject I know is T-L. Ho, PRL **75**, 1186 (1995).

The 331 state is the exact analog of the  $A$  phase of liquid  $^3\text{He}$  (which has no net Cooper pair polarization).

In terms of the original electron coordinates,  $\Psi_{\text{MR}}$  and  $\Psi_{331}$  look totally different, and in particular appear to have a different topology. But when expressed in terms of the composite fermions (see Ho, ref. cit.), it turns out that the only difference is in the spin wave function  $\chi_{\mu\nu}$ !

$$\chi_{\text{MR}} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \chi_{331} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (10)$$

Ho exploits this fact to show that  $\Psi_{331}$  can be deformed continuously into  $\Psi_{\text{MR}}$  without changing “total”  $g(\mathbf{r}_{12}) \equiv \langle \rho(\mathbf{r}_1)\rho(\mathbf{r}_2) \rangle$ , hence without changing  $V_c$ . (Cf.: non-metastability of circulating state of spin-1/2 BEC in annulus).

## 2. The “anti-Pfaffian” state.<sup>8</sup>

If one could neglect completely LL mixing, particle-hole conjugation is an exact symmetry for a half-filled LL. But  $\Psi_{\text{MR}}$  breaks this symmetry, since it is the exact GS of a Hamiltonian which is *not* particle-hole symmetric. So there must exist an “anti-Pfaffian” state ( $\Psi_{\text{AP}}$ ) which is the particle-hole conjugate of  $\Psi_{\text{MR}}$ . For exact particle-hole symmetry it must be degenerate with  $\Psi_{\text{MR}}$ , but LL mixing could stabilize either  $\Psi_{\text{AP}}$  (or  $\Psi_{\text{MR}}$ ). To the best of my knowledge, no-one has formulated the AP wavefunction explicitly in terms of composite fermion coordinates, so one cannot immediately compare it with the MR or 331 states. However, studies of the edge states using bosonization predict different values for the (“universal”) thermal conductance, etc. (see below).

## 3. Other possible identifications.

In the literature there have been yet other suggestions for the nature of the  $\nu = 5/2$  QH state: the “ $K = 8$ ” state, the “ $U(1) \times SU(2)$ ” state and others. I will not go into the details here

Let’s now turn to the important but rather confusing issue of the charge and statistics of the MR states and its competitors. I will try to give a plausible argument only for the MR state, and just quote the results for the others.

For a first pass, let us look at the elementary excitations of a  $(p + ip)$  2D Fermi superfluid. These are (mostly, cf. lecture 27) of two types: simple fermionic BCS quasiparticles, with a minimum excitation energy equal to the gap  $\Delta_0$ , and vortex-antivortex pairs, whose characteristic energy is strongly temperature-dependent as discussed in lecture 10. Now for a strictly 2D system with  $\Delta \sim \epsilon_F$ , the fraction of excited quasiparticles at  $T_{\text{KT}}$  is fairly small (Problem), so let us focus on the vortex-antivortex pairs. Suppose we want to create a vortex at the origin. In a BCS superfluid the way to do this is to simply to multiply the wave function by a factor of the form  $\prod_{i=1}^N f(|z_i|) \exp(i\phi_i/2)$ ,

<sup>8</sup>Levin et al., PRL **99**, 236806 (2007); Lee et al., ibid. 236807.

where  $\phi_i \equiv \arg z_i$  and  $f(|z_i|)$  is some function which tends to zero as  $|z_i| \rightarrow 0$ . The factor of  $1/2$  in the phase corresponds to the well-known fact that in a neutral superfluid such as  $^3\text{He}$  the vorticity is quantized in units of  $h/m_p$  rather than  $h/m$ , where  $m_p \equiv 2m$  is the mass of a Cooper pair. The simplest form of  $f(|z_i|)$  which preserves the analyticity of the wave function up to a cut is  $f(|z_i|) = |z_i|$ ; thus, a vortex at the origin might be created simply by multiplying the groundstate wave function by  $\prod_{i=1}^N z_i^{1/2}$ , and correspondingly a vortex at the position specified by the complex variable  $x + iy \equiv \eta_0$  would then be created by application of  $\prod_{i=1}^N (z_i - \eta_0)^{1/2}$ . Arguing along these lines, we would conclude that a plausible ansatz for a single quasiparticle (actually quasihole) in a system whose GS is described by the MR wave function is

$$\Psi_{\text{qh}} = \prod_{i=1}^N (z_i - \eta_0)^{1/2} \Psi_{\text{MR}}\{z_i\} \quad (11)$$

Whether (11) is correct or not<sup>9</sup> may be a matter of theology, since it actually turns out that it is impossible to create a single isolated vortex in a  $(p + ip)$  Fermi superfluid; for essentially topological reasons one must have either an even number of vortices, or an edge state which plays the same role as a vortex. So the physically meaningful question in the case of the  $\nu = 5/2$  QH state is, what is the correct form of the wave function for *two* quasiholes? The generally accepted ansatz, which preserves all the required symmetries, is

$$\Psi_{2\text{qh}} = \Psi_N^{(\text{L})} \text{Pf} \left\{ \frac{(z_i - \eta_1)(z_j - \eta_2) + (z_i - \eta_2)(z_j - \eta_1)}{z_i - z_j} \right\} \quad (12)$$

where  $\Psi_N^{(\text{L})}$  is the ‘‘Laughlin’’ wave function for  $\nu = 1/2$ , i.e. up to normalization

$$\Psi_N^{(\text{L})} \equiv \prod_{ij} (z_i - z_j)^2 \exp - \sum_i |z_i|^2 / 4l_M^2 \quad (13)$$

Note that this state *cannot* in general be expressed in the form  $f(z_i, z_j) \times \Psi_{\text{GS}}$ . Consider now a 4-qh state (i.e. 2 qh pairs).<sup>10</sup> Start with  $N = 2$ , then a possible wave function. is

$$\Psi_{4\text{qh}} = \Psi^{(\text{L})}(z_1 - z_2)^{-1} \times \{(z_1 - \eta_1)(z_1 - \eta_2)(z_2 - \eta_3)(z_2 - \eta_4) + (z_1 \leftrightarrow z_2)\} \equiv (12)(34) \quad (14)$$

But at first sight there are two other possibilities, namely (13)(24) and (14)(23). However, we now note the identity

$$(12)(34) - (13)(24) = (z_1 - z_2)^2 (\eta_1 - \eta_4)(\eta_2 - \eta_3) \quad (15)$$

from which it follows that

$$(12)(34)(\eta_1 - \eta_2)(\eta_3 - \eta_4) + (13)(24)(\eta_1 - \eta_3)(\eta_2 - \eta_4) + (14)(23)(\eta_1 - \eta_4)(\eta_2 - \eta_3) = 0 \quad (16)$$

<sup>9</sup>In the literature it is usually stated that the single quasihole creation operator is  $\prod_{i=1}^N (z_i - \eta_0)$  as for Laughlin states, but it is not clear that this statement has any real meaning.

<sup>10</sup>Nayak and Wilczek, Nuc. Phys. B **479**, 529 (1996).

i.e. only 2 linearly independent functions. This result also holds for the generalization for  $N > 2$ :

$$\Psi_{4\text{qh}} = \Psi_N^{(L)} \text{Pf} \left\{ \frac{(13)(24)}{z_1 - z_2} - \frac{(13)(24)}{z_3 - z_4} \dots \right\} \quad (17)$$

and it can be generalized to the case of  $2n$  quasiholes :

for  $2n$  quasiholes,  $2^{n-1}$  linearly independent states

NW exhibit an explicit form of a possible choice of basis states:

$$\begin{aligned} \Psi_{4\text{qh}}^{(0)} &= \frac{(\eta_{13}\eta_{24})^{1/4}}{(1 + \sqrt{1-x})^{1/2}} (\Psi_{(13)(24)} + \sqrt{1-x}\Psi_{(14)(23)}) \\ \Psi_{4\text{qh}}^{(1/2)} &= \frac{(\eta_{13}\eta_{24})^{1/4}}{(1 - \sqrt{1-x})^{1/2}} (\Psi_{(13)(24)} - \sqrt{1-x}\Psi_{(14)(23)}) \end{aligned} \quad (18)$$

(?)  $x \equiv \frac{\eta_{12}\eta_{34}}{\eta_{13}\eta_{24}} \quad \eta_{12} \equiv \eta_1 - \eta_2, \quad \text{etc.}$

Let's take  $|x| \ll 1$  i.e.



so that

$$\begin{aligned} \Psi_{4\text{qh}}^{(0)} &= 2^{-1/2}(\eta_{13}\eta_{24})^{1/4} (\Psi_{(13)(24)} + \Psi_{(14)(23)}) \\ \Psi_{4\text{qh}}^{(1/2)} &= 2^{-1/2}(\eta_{13}\eta_{24})^{1/4} (\Psi_{(13)(24)} - \Psi_{(14)(23)}) \end{aligned} \quad (19)$$

Then it is clear that interchange of 1 and 3 (or 2 and 4) affects only the prefactor, so gives a phase factor  $\exp i\pi/4$ . It would thus be natural to take the charge  $e^*$  of a quasihole to be  $e/4$ .

However, interchange of (e.g.) 2 and 3 gives a nontrivial rotation<sup>11</sup> in the space of  $\Psi_0^{(0)}$  and  $\Psi_0^{(1/2)}$ . Thus, the states  $\Psi_{4\text{qh}}^{(0)}$  and  $\Psi_{4\text{qh}}^{(1/2)}$  can in principle be used as the basis for a qubit. More generally,

$$2n \text{ anyons} \rightarrow n \text{ qubits}$$

We now turn to the question: How do we tell experimentally whether the observed  $\nu = 5/2$  QH state is indeed the MR state (as numerical studies tend to suggest) or is one of the competing states ((311), antiPfaffian etc.)? It turns out that all the suggested identifications predict<sup>12</sup> that the effective charge  $e^*$  is  $e/4$ , but they predict different values for the ‘‘Coulomb exponent’’  $g$  which controls some of the proper ties associated with edge states, e.g. temperature-dependence of tunnelling characteristics.

<sup>11</sup>Confirmed by numerical calculations: Tserkovnyak and Simon, PRL **90**, 016802 (2003).

<sup>12</sup>I suspect this follows from rather general topological considerations, see below.



Ansatz	$e^*$	$g$	abelian/nonabelian
MR	$e/4$	0.25	nonabelian
AP	$e/4$	0.5	nonabelian
331	$e/4$	0.375	abelian (?)
$K = 8$	$e/4$	0.125	abelian (?)


**Recent experiments designed to identify the nature of the  $\nu = 5/2$  FQHE state**

1. Dolev et al., (Weizmann Institute) Nature **452**, 829 (2008). Sample: GaAs-AlGaAs heterostructure.

$$n \sim 3 \times 10^{11} \text{ cm}^{-2}$$

$$\mu \sim 3 \times 10^7 \text{ cm}^2/\text{V sec}$$

$$T \sim 10 \text{ mK}$$




Measure: shot noise associated with tunnelling across QPC (quantum point contact). Theoretical prediction:

$$S_I = 2e^*V\Delta g_i t_i(1-t_i) \left[ \coth \left( \frac{e^*V}{2k_B T} \right) - \frac{2k_B T}{e^*V} \right] + 4k_B T g \quad (20)$$

Note that: (a) for  $k_B T \gg e^*V$ , the [ ] goes to 0 so no information on  $e^*$ , (b) for  $k_B T \ll e^*V$ ,  $S_I = 2e^*V\Delta g_i t_i(1-t_i)$ , where  $\Delta g_i = (\nu_i - \nu_{i-1})e^2/h$  with  $t_i \equiv$  tunneling over edge between  $i$  and  $i-1 \Rightarrow$  must know  $\Delta g_i, t_i$  (which may depend on  $I_{\text{imp}}$ ).

Typical data: (note theoretical curves need knowledge of  $g_i, t_i$  - taken from  $\nu = 3$  measurements (?)).

Conclusion:  $e^* = e/4$  with small/zero  $e/2$  contamination, no conclusion about  $g$ .




2. Radu et al. (Harvard-MIT-Lucent), Science **320**, 899 (May 2008). Sample: GaAs-AlGaAs heterostructure.

$$n \sim 2 - 6 \times 10^{11} \text{ cm}^{-2}$$

$$\mu \sim 2 \times 10^7 \text{ cm}^2/\text{V sec}$$

$$T \sim 13 - 60 \text{ mK}$$



Measure:  $V_D$  (i.e.  $R_D$ ) and  $V_{xy}$  (i.e.  $R_{xy}$ ) at fixed  $I_{dc}$  and  $V_g$ , infer the tunneling conductance of the QPC by  $g_T = (R_D - R_{xy})/R_{xy}^2$ . Plot  $R_D (\propto g_T + \text{const})$  as a function of  $I_{dc}$ .

Typical data are shown on the graph above. Fit data to (weak-tunneling) expression

$$g_T = AT^{2g-2}F(g, e^*V/k_B T) \quad V \equiv I_{dc}R_{xy} \quad (21)$$


where  $F$  is a known function. Four fitting parameters:  $A, R^\infty, e^*, g$ . Best fit:  $e^* = 0.17, g = 0.35$  i.e. AP or  $U(1) \times SU_2(2)$  ( $e^* = 0.25, g = 0.5$ ). Data barely consistent with (331) state ( $g = 0.375$ ), probably *inconsistent* with MR ( $g = 0.25$ ).

3. Willett et al., (Lucent), PNAS **106**, 8853 (2 June 09). Sample: GaAs-AlGaAs heterostructure.


$$n \sim 4 \times 10^{11} \text{ cm}^{-2}$$

$$\mu \sim 2 - 5 \times 10^7 \text{ cm}^2/\text{V sec}$$

$$T \sim 25 - 150 \text{ mK}$$



Measure: dependence of  $R_L (\equiv V/I)$  on magnetic field  $B$  and gate voltage  $V_g$ . By calibrating with nearby well-understood QHE plateaux ( $\nu = 5/3, 2, 7/3$ ), can infer




effective area  $A$  of the dot as a function of  $V_g$ . Note effect of  $V_g$  is not primarily through charge accumulation on dot, but directly through change of area  $\Rightarrow$  change of enclosed flux. Hence, should be a unique relation between period observed in  $B$  and  $V_g$ .

Conclusion: at low  $T$ , main component is  $e^* = e/4$ , but with an appreciable  $e/2$  component. At higher  $T$ ,  $e/2$  dominates.


Note: A very recent preprint which may be highly relevant to the interpretation of some or all of these experiments is Ofek et al., arXiv:0911.0794.

**What can we say generically about  $\nu = 5/2$ ?**

A. On torus, by generic Wen-Niu argument, groundstate must be at least doubly degenerate.



B. Hall effect in “wide” Corbino-disk geometry: By original Laughlin argument,  $2\phi_0$  of flux must correspond to  $e$  of charge. So minimum “accessible” periodicity of  $F$  in  $\Phi$  is  $2\phi_0$ : e.g. could have the situation as on the figure with single electron making “adiabatic” transition.



But:  $1/2 = 2/4!$  So, equally plausible scenario is shown on the bottom of the page, with 2 electrons making “adiabatic” transition. This would almost certainly gives “noise” corresponding to  $e/4$  in “constricted” Corbino-disk geometry.

So,  $e^* = e/4$  merely indicates “pairing” and nothing more specific?

[If time permits I will briefly discuss also the  $\nu = 12/5$  state, one candidate state which has anyon excitations of the Fibonacci rather than Ising type.]

