

### Problem Sheet 4

#### 1. ‘Meissner’ and ‘Pauli’ upper critical fields.

Consider a BCS superconductor with no magnetic impurities or spin-orbit coupling, at a temperature  $T$  near  $T_c$  in an external field  $\mathbf{H}$ .

(a) *‘Meissner’ upper critical field.*

For this part of the problem, ignore the coupling of the electron spins to the field and use the GL formalism, ignoring the fourth-order term on the grounds that it will not affect the existence or not of a solution. Show that the condition for a nonzero order parameter to be thermodynamically stable is

$$H < \frac{\Phi_0}{2\pi\xi^2(T)} \equiv H_{c2}^{\text{Meissner}} \quad (1)$$

where  $\xi(T)$  is the GL healing length.

[ Hint: Either use known results on the QHE, etc. (but watch factors of 2!) or use the ‘radial gauge’  $\mathbf{A}(\mathbf{r}) = \frac{1}{2}(\mathbf{r} \times \mathbf{B})$  and the fact that the lowest eigenvalue  $\lambda_n$  of the equation

$$-\frac{1}{r} \frac{d}{dr} r \frac{df}{dr} + \frac{1}{r^2} (n - r^2/2)^2 f = \lambda_n f$$
$$(n = 0, 1, 2, 3 \dots)$$

is 1 independently of  $n$ .]

(b) *‘Pauli’ critical field.*

For this part, ignore the orbital coupling treated in part (a) and consider only the ‘Zeeman’ coupling to the spins. An order-of-magnitude estimate of the largest field which the superconducting state can tolerate in the presence of this interaction is obtained by equating the superconducting condensation energy in zero field to the loss of polarization energy due to formation of (singlet) pairs; to estimate the latter it is

adequate to neglect the nonlinearity of the susceptibility. Show that in this way we obtain for  $T \rightarrow T_c$  the result

$$H < \frac{k_B T_c}{\mu_B} [A'(1 - T/T_c)]^{1/2} \equiv H_{c2}^{\text{Pauli}} \quad (2)$$

and calculate the constant  $A'$ . Compare the latter with the ‘true’ value  $A = 4\pi^2/7\zeta(3)$  obtained from a calculation of the instability of the normal phase, and comment briefly on the reason for any discrepancy.

[ Note:  $1 - Y(T) \rightarrow 2(1 - T/T_c)$  for  $T \rightarrow T_c$  where  $Y(T)$  is the Yosida function. ]

(c) Assume now that for the purpose of order-of-magnitude estimates the formulae obtained for  $H_c$  in parts (a) and (b) can be extrapolated to arbitrary values of  $T/T_c$ . Are ‘Pauli’ effects ever important in the limit  $T \rightarrow T_c$ ? Are they likely to be important for any  $T$  for

- (i) a clean BCS superconductor
- (ii) a very dirty BCS superconductor
- (iii) heavy-fermion systems ( $T_c \sim 1\text{K}$ ,  $H_{c2} \sim 1 - 10\text{T}$ )
- (iv) cuprates ( $T_c \sim 100\text{K}$ , c-axis  $H_{c2} \sim 100\text{T}$ )

(d) Generalize the result of part (a) to the case where the coefficients  $\gamma$ , and hence the healing lengths, are different for the two directions perpendicular to the field. Assuming that the Pauli effect remains isotropic, estimate whether it is likely to be important for fields on the cuprates parallel to the ab-plane (estimated  $H_{c2}$  for this orientation  $\sim 10^3\text{T}$ .)

## 2. ‘Toy model’ to illustrate some aspects of the BdG equations.

Consider the Hamiltonian

$$\hat{H} = (\lambda a_1^\dagger a_2 - i\mu a_1^\dagger a_2^\dagger) + \text{h.c.} \quad (3)$$

where  $a_i$ ’s are fermion operators with the standard anticommutation relations, and the parameters  $\lambda$ ,  $\mu$  are real. Evidently the relevant Hilbert space is 4D and spanned by the vectors  $|n_1, n_2\rangle$ ,  $n_1, n_2 = 0, 1$ . Note that  $\hat{H}$ , while not conserving the quantity  $\hat{n}_1 + \hat{n}_2$ , does conserve its parity (i.e. its ‘evenness’ or ‘oddness’).

- (a) By considering the quantity  $\hat{H}^2$ , or otherwise, find the eigenvalues of  $\hat{H}$ . Is there ever any degeneracy?
- (b) Find the even-parity eigenstates of  $\hat{H}$  for  $|\mu| \geq |\lambda|$  explicitly as linear combinations of  $|00\rangle$  and  $|11\rangle$ , and express them in the form  $(\alpha + \beta a_1^\dagger a_2^\dagger)|00\rangle$ . What are their relative energies?
- (c) Find two linear combinations of the  $a$ 's and  $a^\dagger$ 's which annihilate the even-parity ground state, and two which create from it (normalized) odd-parity states.
- (d) Now write the Bogoliubov quasiparticle creation operator  $\gamma_n^\dagger$  in the form  $\gamma_n^\dagger = \sum_{i=1,2} (u_i^{(n)} a_i^\dagger + v_i^{(n)} a_i)$ . By demanding that  $[\hat{H}, \gamma_n^\dagger] = E_n \gamma_n^\dagger$ , or otherwise, derive the 'BdG' equations and solve for the  $u_i$ 's and  $v_i$ 's for each  $n$ .
- (e) In the special case  $\lambda = \mu$ , show that one of the odd-parity states is degenerate with the even-parity groundstate. By switching to the basis  $|+\rangle \equiv 2^{-1/2}(|1\rangle + |2\rangle)$ ,  $|-\rangle \equiv 2^{-1/2}(|1\rangle - |2\rangle)$ , or otherwise, interpret this result physically.
- (f) How are the results of part (d) related to those of parts (a) and (c)?

### 3. Anomalous (' $\pi$ ') Josephson junction

Consider a tunnel-oxide junction containing magnetic impurities: for simplicity assume them to be polarized at random in the  $\pm z$  directions. Then the transmission matrix element  $T_{\mathbf{kq}\sigma}$  may in general depend on  $\sigma$ : let us write

$$T_{\mathbf{kq}\sigma} = A_{\mathbf{kq}} + \sigma B_{\mathbf{kq}} \quad (4)$$

where  $A$  and  $B$  are assumed to satisfy  $A_{\mathbf{kq}} = A_{-\mathbf{k}-\mathbf{q}}^*$ ,  $B_{\mathbf{kq}} = B_{-\mathbf{k}-\mathbf{q}}^*$  and the quantity  $\overline{A^*B}$  is zero. Assume that the two bulk superconductors connected by the junction are of simple BCS type with  $s$ -wave pairing and that  $T = 0$ .

- (a) Rederive the expression for the Josephson coupling in the form

$$E_J = -\frac{I_c \Phi_0}{2\pi} \cos \Delta\phi \quad (5)$$

and show that under suitable circumstances (what are they?) the quantity  $I_c$  can be *negative*. What relation, if any, can you now obtain between  $I_c$  and the normal-state junction resistance?

- (b) Consider an “rf SQUID” device in which a single such junction, with a negative value of  $I_c$ , is inserted in a bulk superconducting ring (thickness  $\gg \lambda$ ). Show that the total energy of the ring is, including self-inductance effects, has the following form as a function of the total trapped flux  $\Phi$  (which in general includes the contribution  $LI$ ,  $L$  = self-inductance):

$$E(\Phi) = +\frac{|I_c|\Phi_o}{2\pi} \cos 2\pi\Phi|\Phi_o + (\Phi - \Phi_{ext})^2/2L \quad \Phi_{ext} = \text{externally applied flux.} \quad (6)$$

Hence show that for  $L \rightarrow \infty$  there are two degenerate groundstates related by time reversal. What is the energy barrier between them?

- (c) Now consider the effect of the finite self-inductance  $L$  of the ring (but assume zero external flux). Show that below a threshold value  $L_c$ , of  $L$  which depends on  $|I_c|$  the degeneracy is removed, and find  $L_c$ . Find an expression for the height of the barrier for  $L$  just above  $L_c$ .
- (d) Consider specifically a ring with self-inductance 0.1 nH, junction critical current  $|I_c| = 4.4 \mu\text{A}$  and junction capacitance 25 pF. Make a rough estimate of the rate of barrier crossing by thermal activation at (i) 100 mK, (b) 10 mK. Using the result that for a quartic barrier and no ‘detuning’ by external noise, etc., the oscillation rate by quantum tunneling is of order  $\omega_0 \exp -(16V_0/3\hbar\omega_0)$  where  $\omega_0$  is the small-oscillation frequency, estimate this rate and the temperature below which it exceeds the rate of crossing by thermal activation.

[Such a device is contemplated as a possible ‘qubit’]

---

Solutions to be put in 598SC homework box (2nd floor Loomis) by 1 p.m. on Mon. 26 Oct.